CUMULATIVE INDEX 1997

Volume 15

February	TRAVEL RELATED EMERGENCIES, pages 1–281
May	PEARLS, PITFALLS, AND UPDATES, pages 283–490
August	NEUROLOGIC EMERGENCIES, pages 491–733
November	THE USE OF ULTRASOUND IN THE EMERGENCY DEPARTMENT, pages 735–1014

Note: Page numbers of article titles are in **boldface** type

AAA. See *Abdominal aortic aneurysm (AAA).*
ABCs, severe traumatic brain injury, 588–592
Abdominal aortic aneurysm (AAA), aorta in, measurements, 858
  ultrasound examination, 852–856
  clinical presentation and physical examination, 850
  defined, 856
  diagnosis, 851–852
  infrarenal, 857
  natural history, 850
  ruptured, 849–850
    misdiagnosis, 405
  ultrasound findings, 856–858
Abdominal pain, acute abdomen management, 405
  ectopic pregnancy and, 389
  in post-travel patient, 268
Abdominal ultrasound, **825–843**
  in appendicitis, 836–838
  in ascites, 838–839
  biliary tract, 825–834
  liver, 825–826, 834–836
  obstructed bowel, 839–840
  pediatric considerations, 840
    hypertrophic pyloric stenosis, 840–843
    intussusception, 842–843
Abducens nerve (cranial nerve VI), diplopia and, 654, 656
  in neurologic examination, 496–497
Abduction (ocular), 650, 662
Abortion, ultrasound evaluation in first trimester, incomplete, 811
  threatened, 811
Abortive therapy, migraine treatment in children, 630–633
Abruptio placentae, in pregnant traveler, 97
Abscess, of the brain, neuroimaging, 518, 519
  tubo-ovarian, sonographic findings, 817, 818
Accelerated idioventricular rhythm (AIVR), 348
Accessory nerve (cranial nerve XI), in neurologic examination, 499
Accidents, at altitude, 206
Acclimatization, to high altitude, circulatory changes, 193–194
  exercise, 195
  existing medical conditions and, 207–208
  hematologic changes, 194
  high altitude defined, 192–193
  sleep disturbances, 194–195
  special situations, 212
  tissue changes, 194
  unacclimatized rescuers and, 213
  ventilatory changes, 193
Accommodation errors, diplopia and, 655
  headaches associated with, 627
Acetaminophen, for pediatric migraine, 631
Acetaminophen poisoning, *N*-acetylcysteine therapy for, 441–442
Acetazolamide, for acute mountain sickness, 197, 200, 201, 213
N-acetylcysteine (NAC) therapy, 441–442
Achilles reflexes, 501

Achy limb pain, as DCS manifestation, 230
Ackee, toxins from, 173–174
ACLS. See *Advanced cardiac life support (ACLS) protocols.*
Acoustic enhancement, ultrasound images, 955, 959–960
Acoustic nerve (cranial nerve VIII), in neurologic examination, 498
Acoustic shadowing, ultrasound images, 944, 952–955, 956–958
  foreign body detection, 897, 899–900
  gallstones, 953, 955–959
  from splenic flexure, 957
Acoustic window, ultrasound image, 960
  artificial, during scanning, 972
  gallbladder and biliary tract visualization, 826
ACPE. See *Acute cardiogenic pulmonary edema (ACPE).*
Acquired immunodeficiency syndrome (AIDS), CNS infections in patient with, neuroimaging studies, 520–521, 523–525
Active rewarming, of hypothermic patient, core, 470–472
  external, 470
Acute abdomen, pain management in, 405
  in post-travel patient, 269–270
Acute cardiogenic pulmonary edema (ACPE), bi-PAP versus CPAP for, 321–322
  immediate nitrate administration techniques for, 349
Acute mountain sickness (AMS), 191, 452–453
  acetazolamide use in, 200, 201
  dexamethasone use in, 200, 201
  diagnosis, 196
  differential diagnosis, 196
  incidence, 195
  insomnia treatment with or without, 200–201
  natural history, 197
  pathophysiology, 452
  prevention, 198–199
  prevention and treatment, 452–453
  treatment, 197–198, 199
Acute myocardial infarction (AMI), with bundle branch block, diagnosis, 344–345
  cholesterol values following, 348
  diagnosis of, pacemaker and, 342
  thrombolytic therapy versus percutaneous transluminal coronary angioplasty, 354–357
Acute respiratory distress syndrome (ARDS), ventilatory management in patients with, 318
Adamkiewicz, artery of, 700
Adduction (ocular), 650, 663
Adenosine, dysrhythmias and, 345
Admission to hospital. See *Hospital admission.*
Adnexal mass, ectopic pregnancy and, 805, 807, 808
Adult polycystic kidney disease, ultrasound image, 888
Advanced cardiac life support (ACLS) protocols, and cessation of resuscitation in prehospital setting, 297–298
  hemodynamically compromised patient and, 430–432
Adverse drug reactions, recognizing, 437–438
  among current medications, 437
  in elderly, 437–438
  recommendations, 438
AGE. See *Arterial gas embolism (AGE).*
AIDS. See *Acquired immunodeficiency syndrome (AIDS).*
Air-containing structures, ultrasound, 922–923
Air embolism, as diving-related emergency, 234–237
  etiology, 234
  patient transfer, 237
  signs and symptoms, 234–235
  treatment, 236–237
Air travel. See *Commercial air travel.*
Airway management, in prehospital setting, 283–285
  in severe traumatic brain injury, 588–590
Airway rewarming, hypothermic patient, 471
AIVR. See *Accelerated idioventricular rhythm (AIVR).*
AK295, in CNS resuscitation, 529
Alcohol, hypothermia and, 472
Allergic reaction, to anesthesia, 419
ALS. See *Amyotrophic lateral sclerosis (ALS).*
Altitude, 20–23. See also *High-altitude medicine.*
  air travel and, 23, 210
  cardiovascular effects, 20–21
  contact lens use at, 209
  contraindications, 22, 208
  heart disease and, 209
  hypertension and, 209
  information sources, 221–222
  lung disease and, 208–209
  moderate, limitations for, 23
  and pregnancy, 107, 209–210
  pulmonary effects, 21–23
  sickle cell disease and, 209
Altitude sickness, 191, 195
  acute form, 195–201
  cerebral edema, 201–202
  chronic form, 207
  defined, 191–192

"Golden Rules," 211
preexisting medical conditions and, 207–210
pulmonary edema, 202–204
subacute form, 207
treatment for pregnant traveler, 103
other syndromes, 204–206
Amaurosis fugax, 555
Amblyopia, 663
American College of Cardiology, ultrasound training and credentialling guidelines, 914
American College of Emergency Physicians (ACEP), ultrasound training and credentialling guidelines, 913–914
American College of Obstetrics and Gynecology (ACOG), ultrasound training and credentialling guidelines, 914
American College of Rheumatology, fibromyalgia classification criteria, 620
American Institute of Ultrasound in Medicine (AIUM), training and credentialling guidelines, 914
American Spinal Injury Association (ASIA), motor function criteria, 702
AMI. See *Acute myocardial infarction (AMI).*
Amino acid antagonists, excitatory, CNS resuscitation and, 529–533
21-Aminosteroid compounds, in CNS resuscitation, 534
Amiodarone, update on use of, 350–352
Amitriptyline prophylaxis, for pediatric migraine, 633
Amnesia, transient global, 646
Amnestic shellfish poisoning (ASP), 163, 172
Amobarbital interview, 505
A-mode ultrasound, B-mode scan versus, 927
Amoeba, causing traveler's diarrhea, 183
AMPA/KA receptor, 532
in stroke pathophysiology, 554
Amplitude, ultrasound imaging, gain and, 942–944
sound waves, 924, 925
AMS. See *Acute mountain sickness (AMS).*
Amylase, in ectopic pregnancy diagnosis, 391
Amyloidosis, 348
Amyotrophic lateral sclerosis (ALS), 606
Analgesia, oligoanalgesia, 399–400
patient-controlled, for vaso-occlusive crisis, 401–402
for pediatric migraine, 631
for pediatric patient, 385–386, 410–412
for pregnant traveler, 102
for severe traumatic brain injury patient, 598
for stroke patient, 558
Anaphylaxis, travel vaccine reaction, 262
Anechoic structure, on ultrasound image, 941
Anesthesia, in wound care, 417–419
Aneurysm, abdominal aortic. See *Abdominal aortic aneurysm (AAA).*
Angiography, cerebral, in nontraumatic CNS dysfunction, 510
in vascular dissection evaluation, 691–692
Angry patient, how to handle, 306–308
Ankle injuries, evaluation rules for, 375
mortise fractures, failure to recognize, 366, 369
Annular array, ultrasound probe, 934
Anopheles mosquitoes, protection against, 100, 143, 145
Anoxic encephalopathy, as poisoning complication, 436
Anterior cord syndrome, 374, 704
Anterior shoulder dislocations, 365–366
Anterior tibial veins (ATV), in ultrasound examination of lower extremity, 867
Antiarrhythmics, amiodarone, 350–352
proarrhythmias from, 349
Antibiotic drops, for ophthalmologic problems, 332–333
Antibiotics, causing long QT arrhythmias, 349
and digoxin toxicity, 349
for ophthalmologic problems, 332–333
for otitis media, 327–328
for outpatient therapy of community-acquired pneumonia, 320–321
for traveler's diarrhea, 187–188
in pregnant traveler, 109
prophylactic administration, 187
Anticholinergic toxicity, in serotonin syndrome differential diagnosis, 672
Anticonvulsants, rectal administration in pediatric patient, 385
Antidepressants, toxicology, 442–443
Antidiarrheal agents, use by pregnant traveler, 102–103
Antidotes, use in poisoned patients, 443–445
Antiemetics, for pediatric migraine, 631
use by pregnant traveler, 102–103
Antifungal agents, causing long QT arrhythmias, 349
Antigen-binding fragment antidotes, 445
Antihypertensive medications, for stroke patient, 558
Antimalarial drugs, parasitic resistance to, 113–114
Antimotility drugs, for traveler's diarrhea, 185, 187
Antioxidants, in CNS resuscitation, 533–534

Antivenin, for venomous snakebites, 475–476
Anxiety, from flying, 258
Anxiety disorders, weakness associated with, 619
Aorta, abdominal, aneurysm of. See *Abdominal aortic aneurysm (AAA).*
  measurements, 858
  ultrasound examination, 852–856
Aortic dissection, evaluation, 360
Apical window, in 2-D echocardiography, 747, 750
Appendicitis, ultrasound diagnosis, 836–838
Appendix, ultrasound image, 837–838
  inflamed, 838
  normal, 838
ARDS. See *Acute respiratory distress syndrome (ARDS).*
Arrhythmias, amiodarone therapy for, 350–352
  and digoxin toxicity, 348
  long QT, quinine causing, 349
  most commonly missed, 347
  positional, 348
  reperfusion, 348
  VT/wide QRS stable, noninvasive diagnosis, 341–342
Arterial blood supply, to brain, 551–553
  in stroke, 555–557
Arterial gas embolism (AGE), 459–460
Arteries, ultrasound scanning, localization procedure, 907
  for vascular disease diagnosis, 741–742
  visualization technique, 907–908
Arthritis, septic, in pediatric patient, 374–375
Artifacts, ultrasound images, 961
  beam-width, 961–962
  comet tail. See *Comet tail artifact.*
  common types, 968
  contact, 965, 969
  gain, 965, 967
  mirror effect, 965, 967
  reverberation, 963–964
    in foreign body detection, 897
  side lobe, 962–963
Ascites, ultrasound diagnosis, 838–839
Aseptic meningitis, neuroimaging in, 519
ASIA. See *American Spinal Injury Association (ASIA).*
ASP. See *Amnestic shellfish poisoning (ASP).*
Aspiration, ultrasound guidance for, suprapubic, 908
  technique, 909, 910
Aspiration pneumonitis, as poisoning complication, 435
Aspirin, for stroke patient, 558–559
Asthma, external chest compression in, 317
  hypercarbia in, too rapid correction, 318–319
  ipratropium bromide for, 315–316
  oral corticosteroids for, 316
  underdosing of β-agonists in, 319
Ataxia, 504
Atmospheric pressure, 192
Atrial flutter, 347
Atrial septal defect, electrocardiographic identification, 350
Atrioventricular block, response to drugs, 349
Atropine, atrioventricular block response to, 349
Atrovent. See *Ipratropium bromide.*
ATV. See *Anterior tibial veins (ATV).*
Automated implanted defibrillators, interactions with pacemakers, 343–344
  missensing, 343
  oversensing, 343
  pacemaker malfunction after shock, 343–344
  special situations, 344
Automobile crashes, emergency medical services vehicle, 292
Axial resolution, ultrasound images, 944, 946
  beam width on, 948
  pulse dampening on, 951
  pulse duration on, 950
  transducer frequency on, 949

Babinski sign, 501
  in comatose patient, 503
  Bacillus cereus food poisoning, 6, 159
Bacitracin, for suture line aftercare, 424
Bacterial infection, causing traveler's diarrhea, 182
Bag-valve mask, for prehospital airway management, 284
Balance, in neurologic examination, 502
Barbiturates, for severe traumatic brain injury patient, 600
Bare feet, dangers of, 7–8
Barometric pressure, factors affecting, 192
Barotrauma, 457–460
  of ascent, 458–460
  of descent, 457–458
  as diving-related emergency, 23–25
    defined, 226
    ear barotrauma, 226–227
    sinus barotrauma, 227–229
  external ear, 458
  inner ear, 457
  pulmonary, 458–459
Basic fibroblast growth factors (bFGF), in CNS resuscitation, 536–537
Baye's theorem, diagnostic testing and, 304
BDNF. See *Brain-derived neuronotropic factor (BDNF).*

Beam, ultrasound, characteristics, 954
safety issues, 973–974
Beam width, on resolution, ultrasound images, 948
Beam-width artifact, ultrasound images, 961–962
Beclomethasone, for chronic bronchospastics, 316
Bedside echocardiography, characteristics, 736–737
for chest pain evaluation, 758
limitations, 737–739
Beer, CO_2 end-tidal readings and, 348–349
Behaviorally-disturbed patients, COBRA transfer rules, 287
Behavioral therapy, migraine treatment in children, 633
Bell's palsy, 331–332, 498
"Bends," 23–25. See also *Decompression sickness (DCS).*
Benedikt's syndrome, 656–657, 663
β-agonists, synergism with ipratropium bromide, 316
underdosing in asthma, 319
Beta blockers, coronary artery disease patients and, 33–34
Betadine, for wound irrigation, 420
bFGF. See *Basic fibroblast growth factors (bFGF).*
Bilevel positive airway pressure (Bi-PAP), for acute pulmonary edema, 321
Biliary tract, sonographic evaluation, 825–834
acoustic window for visualization, 826
distention, 831–832, 833
pathology, 828–834
Binocular diplopia, 658, 663
Binocular vision, 650, 651
requirements for, 649
Biologic effects, ultrasound images, 972–974
Bi-PAP. See *Bilevel positive airway pressure (Bi-PAP).*
Bismuth subsalicylate, for traveler's diarrhea, 185, 187
prophylactic administration, 187
Bites, weakness induced by, 617
Black structure, on ultrasound image, 941
Bladder irrigation, hypothermic patient, 471
Bland infarcts, 515
Bleeding, vaginal, ectopic pregnancy and, 389
ultrasound evaluation, first trimester, 800–811. See also *Ectopic pregnancy.*
second- and third-trimester, 813–815
Blindness, in conversion disorder, 504
Blink reflex, in comatose patient, 503
Blood, sonographic appearance, in blunt abdominal trauma, 771, 772
Blood-brain barrier, CNS resuscitation and, 533–534
in severe traumatic brain injury, 591
Blood pressure, control in stroke patient, 558
resuscitation in severe traumatic brain injury, 590–592
Blood smear, for malaria diagnosis, 128–132
Blood supply, to brain, 551–553
in severe traumatic brain injury, 585–588
in stroke, 555–557
to spinal cord, 700
Blood tests, for pediatric patient, 382
Bloody diarrhea, in post-travel patient, 268
Blowfish poisoning, 162, 168–170
Blunt trauma, abdominal, ultrasound in, **763–785**
in adults, 774–781
in children, 781–782
diagnostic algorithm, 778–779
equipment, 765–766
evaluation algorithm, 783
history, 763–765
limitations, 774
technique, 766–774
training, 782–785
to chest, transesophageal echocardiography in, 360–361
to head and neck, carotid and vertebral artery dissections, mild injury, 689
serious injury, 690–691
to heart, echocardiographic diagnosis, 761
to kidney, formal ultrasound evaluation, 882–883
B-mode ultrasonography, 926
in abdominal aortic aneurysm diagnosis, 851, 852
A-mode versus, 927
Boating injuries, recreational, 245
Boiling of water, for diarrhea prevention, 184
Bony alignment rules, foot, 369, 370
Botulism, 160–161, 610, 615–616, 657
Bowel pathology, ultrasound diagnosis, intussusception, 842–843
obstruction, 839–840
Boyle's law, 224–225
Brain, anatomy, 551–553
secondary insults in severe traumatic brain injury, 582–584
Brain abscess, neuroimaging, 518, 519
Brain death, **713–721**, 715–716
cardinal findings in, 716–718
complicating factors, 718
confirmatory testing, 718
criteria for, current consensus, 716–720
evolution of, 715–716

Brain death *(Continued)*
and determination of death, historical approaches, 714
documentation, 718
in emergency medicine, 713–714
family consultation in, 719
misdiagnosis, 719–720
in pediatric patient, 719
in potential organ donor patient, 720–721
prerequisites prior to evaluation for, 716
Brain-derived neuronotropic factor (BDNF), in CNS resuscitation, 536–537
Brain hemorrhage, neuroimaging, 511–514, 515
cerebellar, 514
intracerebral, 512–514
intraventricular, 514, 515
subarachnoid, 511, 512
subdural, 511–512
Brain injury, fluid percussion, 528–529
mild head trauma, **563–576.** See also *Mild head trauma.*
severe traumatic, **581–600.** See also *Severe traumatic brain injury.*
Brain tumors, neuroimaging, 523–524
Bronchiolitis, ipratropium bromide for, 315–316
Bronchitis, at high altitude, 205
Bronchospastic patient, discharged, prednisone and, 319
steroid metered-dose inhaler for, 316
MDI-spacer therapy for, 322
post-travel, parasite-induced bronchospasm, 274–275
Brown-Sequard syndrome, 374, 704
Brudzinski's sign, 501
Bruises, evaluation in pediatric patient, 386–387
Bucket-handle fractures, in children, 370, 372
Bullets, ultrasound scanning, 902, 903–905
Bundle branch block, myocardial infarction with, diagnosis, 344–345
Buprenorphine, for renal colic pain management, 406
Buspirone, serotonin syndrome and, 670
Button battery, lodged in pediatric esophagus, 386
Butyl-2-cyanoacrylate, for wound closure, 423
BW 619C89, and glutamate release, 532

Cactus spine foreign body, ultrasound image, 901
CAD. See *Carotid artery dissection (CAD); Coronary artery disease (CAD).*
Calcium channel blockers, CNS resuscitation and, 528–529
as prophylaxis for pediatric migraine, 633
Calf-vein deep venous thrombosis, isolated, diagnostic ultrasound for, 868
Calorimetric end-tidal carbon dioxide detector, disposable, 287
Calpain inhibitors, CNS resuscitation and, 528–529
Campylobacter jejuni, causing traveler's diarrhea, 182
Cancer, oropharyngeal, 329
ultrasound findings, gallbladder, 833–834
liver, 836
CAP. See *Community-acquired pneumonia (CAP).*
Capnometers, endotracheal versus esophageal intubation and, 905–906
Captopril, sublingual, 348
Carbamates, as cause of weakness, 611
Carbon monoxide exposure, 465–469
arterial versus venous measurements, 468
and cyanide poisoning combined, 467
epidemiology, 465
in infants, falsely elevated COHb levels, 467
management, 466
organ transplantation following, 467
prevention, 465
prognostic modalities, 466–467
pulse oximetry, 467
Carcinoma. See *Cancer.*
Cardiac arrest, survivorship data, 285–286
Cardiac disease, air travel and, 256–257
high-altitude travel and, 209
maritime medicine and, 244
Cardiac ischemia, acute, diagnostic ultrasound for, 742
Cardiac motion, fetal, sonographic detection, 799
Cardiac output, and oxygen delivery, effects of poisoning on, 432
Cardiac rupture, echocardiographic diagnosis, 761
Cardiac tamponade, echocardiographic diagnosis, 750–755, 757
Cardiac trauma, echocardiographic diagnosis, 759–761
Cardiac ultrasound, **745–761.** See also *Echocardiography.*
Cardiogenic shock, echocardiographic diagnosis, 757
Cardiology, **341–361**
pearls, automated implanted defibrillators, 343–344
miscellaneous, 348–350
myocardial infarction, diagnosis of, with bundle branch block, 344–345

pacemakers and, 342
QRS rhythms, diagnosis of, regular wide, 345–347
VT/wide stable arrhythmias, 341–342
pitfall, arrhythmias most commonly missed, 347
updates, amiodarone, 350–352
cardiovascular imaging modalities, 352–354
thrombolytic therapy, 357–359
versus percutaneous transluminal coronary angioplasty, 355–357
transesophageal echocardiography, 359–361
Cardiopulmonary resuscitation (CPR), 284
cough technique, 349
in-flight emergency requiring, 256
Cardiorespiratory disorders, and grunting respiration in pediatric patient, 384
Cardiovascular system, altitude and, 20–21
heat tolerance and, 19
imaging, in coronary artery disease diagnosis, 352–354
update on, 352–354
problems, air travel and, 256–257
maritime medicine and, 244
Carotid artery, and vertebral artery dissections, **677–695.** See also *Carotid artery dissection (CAD); Vertebral artery dissections (VAD).*
diagnostic evaluation, 691–692
diagnostic pitfalls and dilemmas, 693–695
intracranial, 688
pathophysiology, 678–680
spontaneous, 680–688
traumatic, 688–691
treatment, 692–693
Carotid artery dissection (CAD), pain distribution in, 681
spontaneous, 680–683
case report, 680
clinical course or prognosis, 683
clinical features, 686
clinical presentation, 680–682
intracranial, 688
predisposing factors and mechanisms of injury, 682–683
warning signs and symptoms, 682
traumatic, 688
mild blunt injury, 689
serious blunt or penetrating injuries, 690–691
Carotid sinus massage, in diagnosis of spells, 645
Carpal dislocations, failure to recognize, 366, 367
Case management, telemedicine services, 246
Cassava, toxins from, 174
CBC. See *Complete blood count (CBC).*
CBF. See *Cerebral blood flow (CBF).*
CBV. See *Cerebral blood volume (CBV).*
CCP. See *Cerebral perfusion pressure (CCP).*
Celiac axis, in ultrasound examination of aorta, 853, 855
Central cord syndrome, 374, 704
Central effects, ultrasound images, 944, 952–960
acoustic enhancement, 955, 959–960
acoustic shadowing, 897, 899–900, 944, 952–955, 956–959
Central nervous system (CNS), dysfunction, neuroimaging in. See *Neuroimaging, in nontraumatic CNS dysfunction.*
peripheral nerve lesions versus CNS disease, 605–606
resuscitation, **527–538.** See also *Resuscitation, central nervous system.*
Central veins, ultrasound examination, 870–871
Central vein thrombosis, diagnostic ultrasound for, 868–869
imaging criteria, 871
imaging studies in evaluation, magnetic resonance imaging, 870
ultrasonography, 869–870
other techniques, 870
Cephalosporin, for streptococcal pharyngitis, 335
Cerebellar hemorrhage, neuroimaging, 514
Cerebral angiography, in nontraumatic CNS dysfunction, 510
Cerebral blood flow (CBF), in severe traumatic brain injury, 585–588
Cerebral blood volume (CBV), in severe traumatic brain injury, 587
Cerebral computed axial tomography, subarachnoid hemorrhage diagnosis, 304
Cerebral edema, diffuse, neuroimaging, 517–518
at high altitude. See *High altitude cerebral edema (HACE).*
Cerebral perfusion pressure (CCP), in severe traumatic brain injury, 585–588
Cerebrovascular accidents. See *Stroke; Stroke syndromes.*
Cerebrovascular disease, 646
ischemic, neuroimaging, 514–517
Cervical spine fractures, 373–374
CFD. See *Color flow Doppler (CFD).*
CHD. See *Coronary heart disease (CHD) patients.*
Chemoprophylaxis, for malaria prevention, 145–148
in pregnant traveler, 100–101
Chest compression, external, in asthma, 317

Chest injuries, blunt and penetrating, transesophageal echocardiography in, 360–361
Chest pain, estradiol drips for, 349
CHF. See *Chronic heart failure (CHF).*
Chickenpox immunization. See *Varicella immunization.*
Children, in emergency department. See *Pediatric patient; Pediatrics.*
travel with. See *Infants and children, travel with.*
Chloral hydrate, pediatric use, 386, 412
Chlorine, for water purification, 184
Chloroquine, for malaria prevention, 100, 145, 146–147
for malaria treatment, 266, 267
"Chocolate cysts," sonographic findings, 819
Cholecystitis, ultrasound findings in, 830–831
acute form, 831, 832
emphysematous form, 958
gangrenous form, 831
Choledocholithiasis, ultrasound findings, 831
Cholelithiasis, ultrasound findings, 828–830. See also *Gallstones.*
Cholera immunization, 53, 59
for elderly, 66
for immunocompromised person, 66
for infants and children, 66, 80
in pregnancy, 66, 99
Cholesterol values, following acute myocardial infarction, 348
Chronic fatigue syndrome, 618, 619
Chronic heart failure (CHF), beta blocker eye drops and, 348
CPAP mask use in, 349
pleural effusions in, 348
Chronic meningitis, neuroimaging in, 520
Chronic mountain sickness (CMS), 207
Chronic obstructive pulmonary disease (COPD), hypercarbia in, too rapid correction, 318–319
ipratropium bromide for, 315–316
oral corticosteroids for, 316
Chronic parvo virus B17 syndrome, weakness associated with, 618
CIE. See *Counterimmunoelectrophoresis (CIE).*
Ciguatera fish poisoning, 162, 165–168
diagnosis, 167
geographic distribution, 166
pathophysiology, 166
in post-travel patient, 272
spectrum of illness, 165–166
toxicology, 166–167
treatment and prevention, 167–168
Ciguatoxin, 617
Circadian rhythm, jet lag and, 11
Circle of Willis, 552–553
Circulatory changes, at high altitude, 193–194
Cirrhosis of the liver, ultrasound image, 943
Cis-atracurium, use in prehospital setting, 299
Claritin, causing long QT arrhythmias, 349
Claude's syndrome, 656, 663
Clindamycin, malaria treatment with, 266
prevention in pregnant traveler, 100
Clinical practice, of emergency medicine, **303–312**
pearls, angry patient, how to handle, 306–308
diagnostic testing, 303–304
Jehovah's Witnesses, 312
minors, treatment of, 310–311
physician-patient interactions, 305–306
physician-physician interactions, 308–309
pitfalls, admission orders, 310
EMTALA, 309–310
minors, and refusal of care, 311–312
tympanic thermometers, 304–305
update, lumbar puncture, 304
Clostridium botulinum toxin, 615–616
Clostridium perfringens food poisoning, 159
Closure, of wounds, 421–424
Clothing, for travelers, 9
Clozaril (clozapine) overdose, 443
CMS. See *Chronic mountain sickness (CMS).*
CNQX. See *6-Cyano-7-nitroquinoxaline-2,3-dine (CNQX).*
Coagulation, at high altitude, 204
Cochlear nerve, in neurologic examination, 498
Coins, lodged in pediatric esophagus, 386
Colchicine, for pericarditis relief, 348
Cold agglutinin test, for *Mycoplasma pneumoniae,* 316–317
Cold caloric testing, in comatose patient, 503
Cold exposure, traveler and, 18–19
at altitude, 206
as HAPE predisposing factor, 202
Colds, treatment for pregnant traveler, 103–104
Colloid resuscitation, in severe traumatic brain injury, 591
Colorado Medical Society, management guidelines for sports concussions, 576
Color-flow Doppler (CFD), ovarian torsion detection by, 820
vascular evaluation, 860
lower extremity deep vein thrombosis, 864–865
Comatose patient, neurologic examination, 502–503
Combitube airway, 284

Comet tail artifact, ultrasound scanning, endotracheal intubation, 906–907
foreign body detection, 897, 898
Comitance, 663
spread, 663
Commercial air travel, cabin pressure, 210
medical emergencies in, **251–259**
anxiety, 258
cardiac disorders, 256–257
demographics, 251–252
dysbarism, 258
ear and sinus disorders, 257
"economy class syndrome," 258
gastrointestinal disorders, 257–258
in-flight death causes, 252
in-flight physiology and pathophysiology, 252–253
medical training and equipment, 253–254
operational procedures and unscheduled landings, 254–255
pulmonary disorders, 255–256
syncope, 255
trauma, 259
medical restrictions to, 23, 210
by pregnant traveler, 28, 107–108, 210
Commercial fishing, medical emergencies and. See *Maritime medicine.*
Common artifacts, ultrasound images, 968
Common bile duct dilatation, ultrasound findings, 832–833
Communication, physician-patient, 305–306
angry patient or family, 306–308
elderly patient, 337–338
physician-physician, 308–309
positive behaviors, 306
Communication logistics, telemedicine services, 247–248
Community-acquired pneumonia (CAP), antibiotic choice for outpatient therapy, 320–321
Compartment syndrome(s), diagnosis, 373
intraosseous infusion-induced, 373
as poisoning complication, 436
thrombolytic-induced, 373
Compazine, for migraine, 408–409
Complete blood count (CBC), for pediatric patient, 382
Complete cord syndrome, 703–704
Complete neurologic examination, directed examination versus, 492
Complicated migraine, 628
Comprehensive Omnibus Budget Reconciliation Act of 1986 (COBRA), patient transfers rules, 287–289
penalties for violating, 288
Compression, of chest, in asthma, 317
during ultrasound scan, 970–971
ultrasound scanning technique, 972
Computed axial tomography, cerebral, subarachnoid hemorrhage diagnosis by, 304
Computed tomography (CT), aorta measurements, 858
in aortic dissection evaluation, 360
in cholelithiasis detection, ultrasound versus, 825
in diagnosis of spells, 644
in diplopia evaluation, 662
in foreign body detection, 896
in minor head injury, indications, 566, 567
low-risk patient identification and, 566–568
studies against, 571–572
studies in favor of, 568–571
summary of studies, 572–573
in nontraumatic CNS dysfunction, 509
contrast media for, 510–511
in pediatric migraine diagnosis, 630
in renal colic diagnosis, 879, 881
in severe traumatic brain injury, prioritization, 596–598
ultrasonography versus, in blunt abdominal trauma evaluation, in adults, 774–781
in children, 782
in vascular dissection diagnosis, 694–695
Concussions, postconcussion syndrome and, 574–575
sports injuries and, 575
management guidelines, 576
Conjugate gaze, 654, 655
Consent, emergency, for minors, 310–311
for transfer, 289
Consultation, physician-physician, 308–309
Contact artifact, 965, 969
Contact lens use, high altitude travel and, 210
Contact scanner, ultrasound probe, 936
Continuous positive airway pressure (CPAP), for acute pulmonary edema, 321
mask use in chronic heart failure, 349
Contrast nephropathy, 510
Contrast venography, in lower extremity deep vein thrombosis evaluation, 860–861
Conus medullaris syndrome, 705
Convergence errors, diplopia and, 655
Conversion disorder, suspected, neurologic examination in, 503–505
weakness associated with, 619
Coordination, in neurologic examination, 502
COPD. See *Chronic obstructive pulmonary disease (COPD).*
Cord syndromes. See *Spinal cord syndromes.*

Corneal abrasions, antibiotic therapy for, 333
  corneal abrasions for, 333
Corneal eyepad, 333
Corneal light reflex test, 661
Corner fractures, in children, 370, 372
Corner sutures, for wound closure, 422–423
Coronary angioplasty, versus thrombolytic therapy for acute myocardial infarction, 355
Coronary artery disease (CAD), cardiovascular imaging modalities in diagnosis of, 352–354
Coronary heart disease (CHD) patients, wilderness travel and, 30–34
  asymptomatic patients, 32–33
  exercise testing, 31
  medication considerations, 33–34
  symptomatic patients, 32
Corpus luteum cysts, sonographic findings, 819
Corticosteroids, oral, for chronic bronchospastics, 316
Cough, at high altitude, 205
  parasite-induced, 274–275
  treatment for pregnant traveler, 103–104
Cough CPR, 349
Council of Residency Directors (CORD), ultrasound curriculum, 915–916
Counterimmunoelectrophoresis (CIE), in pediatric meningitis evaluation, 384
Cover test, 661
CPAP. See *Continuous positive airway pressure (CPAP).*
CPK. See *Creatinine phosphokinase (CPK).*
CPR. See *Cardiopulmonary resuscitation (CPR).*
Cranial nerves, lesions causing diplopia, 656–657, 662
  in neurologic examination, 495–499
    abducens, 496–497
    accessory, 499
    acoustic, 498
    facial, 497–498
    glossopharyngeal, 499
    hypoglossal, 499
    oculomotor, 496–497
    olfactory, 495
    optic, 496
    trigeminal, 497
    trochlear, 496–497
    vagus, 499
  in ocular motor system, 652–654
Crashes, emergency medical services vehicle, 292
Creatine kinase, in ectopic pregnancy diagnosis, 391
Creatinine phosphokinase (CPK), in diagnosis of spells, 643
Credentialling issues, in emergency ultrasound, **913–919**
  approval strategies, 917–919
  qualification pathway, 915–916
  quality assurance, 916–917
  training and guidelines, 914–915
Creeping eruption, in post-travel patients, 277–278
Cricothyroidotomy, following neuromuscular blockade, 298
"Crochetage," 350
Cross-cover test, 661
Croup, racemic epinephrine for, 330–331
Crown-rump length, sonographic measurement, 799
Cryptococcosis, in AIDS patient, neuroimaging studies, 521, 525
Cryptococcus neoformans infection, in AIDS patient, neuroimaging studies, 521, 525
Cryptosporidium, causing traveler's diarrhea, 183
Crystalloid resuscitation, in severe traumatic brain injury, 591
CT. See *Computed tomography (CT).*
Culiseta impatiens mosquitoes, protection against, 7
Curved linear scanner, ultrasound probe, 934, 936
Cutaneous decompression sickness, 456
Cutaneous leishmaniasis, in post-travel patients, 279
Cutaneous myiasis, in post-travel patients, 278
Cyanide poisoning, carbon monoxide exposure combined with, 467
Cyanide toxicity, from cassava, 174
6-Cyano–7-nitroquinoxaline–2,3-dine (CNQX), and NIDA receptor activation, 531–532
Cyclotorsion, 650, 663
Cyproheptadine prophylaxis, for pediatric migraine, 633
Cysts, female pelvis, sonographic findings, 819
  hepatic, ultrasound-guided drainage, 909, 910

Dalton's law, 225
DAN (Divers' Alert Network), 223, 456
DCS. See *Decompression sickness (DCS).*
Death, determination of, historical approaches, 714. See also *Brain death.*
  in-flight, causes, 252
Decadron. See *Dexamethasone.*
Deceleration injury, in mild head trauma, 564
Decision rules, for acute knee injury evaluation, 375–376

for ankle evaluation, 375
Decompression sickness (DCS), 23–25, 455–457
clinical presentations, 456
defined, 229
manifestations, 230–231
symptom onset, 229–230
treatment, 456–457
adjunctive therapies, 234
hyperbaric oxygen therapy, 231–234
vestibular, 231
Decontamination, of wounds, 419–420
DEEP. See *Definite ectopic pregnancy (DEEP).*
Deep femoral vein (DFV), in ultrasound examination of lower extremity, 866
Deep tendon reflexes, 501
and spinal cord level relationship, 702–703
Deep vein thrombosis (DVT), diagnostic ultrasound for, isolated calf-vein, 868
lower extremity, 860, 862–867. See also *Lower extremity deep vein thrombosis (LEDVT).*
possible future use in emergency department, 741
proximal lower extremity, evaluation, 323–324
DEET (N,N,-diethyl-meta-toluamide) insect repellent, 7, 100
against mosquitoes, 143, 145
use by pregnant traveler, 105
Defibrillation, automated implantable devices. See *Automated implanted defibrillators.*
early, emergency medical services and, 285–286
in-flight emergency requiring, 256
Definite ectopic pregnancy (DEEP), 804
defined, 800, 801
Dehydration, maritime medicine and, 244
Delayed intracranial injuries, in mild head trauma, 573
Delirium, clinical characteristics, 494
evaluation in neurologic examination, 494–495
in serotonin syndrome differential diagnosis, 672
Dementia, clinical characteristics, 494
evaluation in neurologic examination, 494–495
Depression (ocular), 650, 663
Dermacentor, envenomation by, 610, 617
Desyrel (trazodone) overdose, 443
Deterioration, of patient, EMTALA and, 309–310
Devastator bullets, ultrasound scanning, 902, 903–905
Dexamethasone, for acute mountain sickness, 197, 200, 201, 213
Dextromethorphan, and NMDA receptor antagonism, 530–531
Dextrorphan, and NMDA receptor antagonism, 530–531
DFV. See *Deep femoral vein (DFV).*
DHE. See *Dihydroergotamine (DHE).*
Diabetic patients, wilderness travel and, 34
Diagnostic peritoneal lavage (DPL), ultrasonography versus, in blunt abdominal trauma evaluation, 764
in adults, 774–781
Diagnostic testing, appropriate use of, 303–304
Diamox. See *Acetazolamide.*
Diaphragm, ultrasound image, 942
Diaphyseal long-bone fractures, nonaccidental, in children, 370, 372
Diarrhea, in post-travel patients, 268–272
acute abdomen, 271–272
bloody diarrhea and abdominal pain, 270
ciguatera fish poisoning, 272
lactose intolerance, 269–270
schistosomiasis, 270–271
traveler's diarrhea, 268–269
typhoid fever, 271
traveler's. See *Traveler's diarrhea.*
Diarrheal shellfish poisoning (DSP), 163, 172
Diarrhea travel kit, 184
Diet, jet lag and, 11
N,N,-Diethyl-meta-toluamide insect repellent. See *DEET (N,N,-diethyl-meta-toluamide) insect repellent.*
Diffuse cerebral edema, neuroimaging, 517–518
Digoxin toxicity, antibiotic administration and, 349
arrhythmias and, 348
in elderly, 349
Dihydroergotamine (DHE), for pediatric migraine, 631
serotonin syndrome and, 670
Dimethyl sulfoxide (DMSO), in CNS resuscitation, 533
Diphtheria, 657
as cause of weakness, 610
Diphtheria immunization, adult, 55
routine, in childhood, 48, 78
special groups recommendations, 66
Diplopia, 496–497
clinical approach to, 659–662
defined, 649
diagnostic tests for, 661
in the Emergency Department, **649–664**
etiology, 655–658
heteronymous, defined, 663
homonymous, defined, 663
intraocular structures and, 649–650
monocular, 658–659

Diplopia *(Continued)*
ocular motor system and, cranial nerves, 652–654
muscles, 650–652, 653
supranuclear control, 654
Directed neurologic examination, versus complete examination, 492
Disease prevalence, diagnostic testing and, 303–304
Disk battery, lodged in pediatric esophagus, 386
Dislocations, anterior shoulder, 365–366
carpal, failure to recognize, 366, 367
failure to recognize, Lisfranc fracture, 369–371
Disposable calorimetric end-tidal carbon dioxide detector, 287
Dissections, carotid and vertebral artery, **677–695.** See also *Carotid artery dissection (CAD); Vertebral artery dissections (VAD).*
diagnostic evaluation, 691–692
spontaneous, 680–688
traumatic, 688–691
Divers' Alert Network (DAN), 223, 456
Diving, injuries associated with. See *Diving-related emergencies.*
medical contraindications, 24
risks associated with, 23–25
Diving-related emergencies, **223–237**
air embolism, 234–237
barotrauma, ear, 226–227
sinus, 227–228
decompression sickness, 228–231
treatment, 231–234
gas laws, 224–226
incidence, 223
pressure changes, 224
Dizocilopine, and NMDA receptor antagonism, 530–531
Dizziness, evaluation in neurologic examination, 492–493
vestibular nerve and, 498
DMSO. See *Dimethyl sulfoxide (DMSO).*
Dobutamine stress echocardiography, in coronary artery disease diagnosis, 354
Documentation, brain death, 718
neurologic examination, 505–506
Doppler effect, 859
Doppler imaging, in vascular dissection evaluation, 692
Doppler shift, 859
Doppler ultrasound, abdominal aortic aneurysm, 857–858
principles, 859–860
blunt renal trauma evaluation, 884
color-flow. See *Color-flow Doppler (CFD).*
continuous-wave, in lower extremity deep vein thrombosis evaluation, 862–863
Doxycycline, pregnant traveler and, diarrhea treatment, 108
malaria prevention, 100, 146, 148
malaria treatment, 266
DPL. See *Diagnostic peritoneal lavage (DPL).*
Drainage, of fluid-filled structures, ultrasound guidance for, 908–909, 910
Drinking. See *Water.*
Driving, international travel and, 6
Drop attacks, 646
Drowning. See *Near drowning.*
Drug-induced weakness, 608, 616
Drug screening, qualitative versus quantitative, 430
urine drug screen, 430
Drug use, hypothermia and, 472
Dry field, for tourniquet examination of extremities, 371
DSP. See *Diarrheal shellfish poisoning (DSP).*
Dysbarism, air travel and, 258
Dysrhythmias, incorrect drug use for, 345

Ear, nose, and throat (ENT), **327–338.** See also *Ear disorders.*
pearls, epistaxis patient, management recommendations for, 336–337
foreign body removal, 328–329
iritis diagnosis, 333–334
patches and medications, for ophthalmologic problems, 332–333
racemic epinephrine, use in croup patients, 330–331
sore throat, 335–336
steroid use, in upper airway disease, 328
pitfalls, elderly, communication difficulties with, 337–338
head and neck malignancies, identification of, 329–330
sore throat, 335–336
updates, Bell's palsy, 331–332
strep throat, treatment for, 334–335
Ear barotrauma, as diving-related emergency, 226–227
Ear disorders, air travel and, 257
wilderness travel and, 36
Eating. See *Food.*
ECG. See *Electrocardiogram (ECG).*
Echocardiography, 745–750
dobutamine stress, in coronary artery disease diagnosis, 354
establishing diagnosis, cardiac tamponade, 750–755
cardiac trauma, 759–761
hypotension of unknown cause, 756–757, 758
ischemic heart disease, 757–758
pulseless electrical activity, 755–756

M mode, 940
transesophageal versus transthoracic, 359–361
windows used during, apical, 747, 750
left parasternal, 746–749
subcostal, 745–746, 747
Echogenic image, ultrasound, 923
Echo patterns, foreign bodies, ultrasound detection technique, 897–901
"Economy class syndrome," 258
Ectopic pregnancy, 800
adnexal mass, 805, 807, 808
definite, 800, 801, 804
diagnosis, 390–391
empty uterus and, 809
extrauterine empty gestational sac and, 804
incidence estimates, 803
interstitial, 808
and intrauterine pregnancy combined, 808–809
methotrexate treatment, 393–394
pseudobicornuate, 809
pseudogestational sac of, 809, 810
rare forms, 809
ruptured, 805, 806–807
ruptured, tachycardia and, 389–390
screening for, 389
serologic markers, 391–392
sonographic diagnosis, pitfalls, 810–811
protocol to rule out, 800–803
sonographic spectrum, 803–810
tubal ring and, 804–805, 806
Edema, acute cardiogenic pulmonary. See *Acute cardiogenic pulmonary edema (ACPE).*
diffuse cerebral, neuroimaging, 517–518
at high altitude. See *High-altitude cerebral edema (HACE); High-altitude pulmonary edema (HAPE).*
Edrophonium challenge test, in myasthenia gravis diagnosis, 614
Education, emergency medical services, medical oversight lacking in, 293–295
telemedicine services, 247
traveler's diarrhea, 247
EEG. See *Electroencephalogram (EEG).*
Effects, ultrasound images, biologic, 972–974
central, 944, 952–960
lateral cystic shadowing, 924, 925, 960–961
Effexor (venlafaxine) overdose, 443
Elderly, adverse drug reactions in, recognizing, 437–438, 439
communicating with, 337–338
digoxin toxicity in, 349
hypothermia in, 472
immunizations prior to travel, 66, 67
wilderness travel and, 30
Electrocardiogram (ECG), in diagnosis of spells, 643–644
Electroencephalogram (EEG), in diagnosis of spells, 644
Electrolyte-induced weakness, 610–611, 616–617
hyperkalemia, 611, 617
hypermagnesemia, 611, 617
hypokalemia, 611, 616–617
hypophosphatemia, 610, 617
Electrolyte levels, in diagnosis of spells, 643
Electronic sector scanner, ultrasound probe, 934, 935
Electrophysiologic testing, in diagnosis of spells, 645
Elevation (ocular), 650, 663
Embolism, during air travel, 258
arterial gas, 459–460
diving-related. See *Air embolism.*
pulmonary. See *Pulmonary embolism.*
Embryonic pole, sonographic measurement, 799
Emergencies. See *Medical emergencies.*
Emergency department, travel medicine problems encountered in, **261–280**
diarrhea, 268–272
malaria, 264–267
respiratory tract infections, 272–275
skin lesions, 277–279
travel vaccine reactions, 261–264
worms and parasites, 275–277
ultrasound in, applications of, **735–743**
examination characteristics, 736–737
future applications, 740–743
limitations, 737–739
scope, 739–740
vascular evaluation, 859
Emergency medical services (EMS), and early defibrillation, 285–286
education and practice, medical oversight lacking in, 293–295
Emergency medical technician-basic (EMT-B) curriculum, airway management, 283
defibrillation, 283
update on, 295–296
Emergency medical technician (EMT), airway management by, 283–285
confirming correct placement of endotracheal intubation, 286–287
early defibrillation by, 285–286
fluid resuscitation of trauma patients in prehospital setting, 296–297
neuromuscular blockade administered by, 298–299
Emergency Medical Treatment and Active Labor Act (EMTALA), patient transfer rules, 288–289
screening examination requirements, 309–310

Emergency medicine, clinical practice, **303–312.** See also *Clinical practice, of emergency medicine.*
Emergency physicians. See *Physicians.*
Emergency ultrasound, credentialling issues in, **913–919**
qualification pathway, 915–916
quality assurance, 916–917
strategies, 917–919
training and guidelines, 914–915
Emphysematous cholecystitis, ultrasound image, 958
Empyema, neuroimaging, 518–519
EMS. See *Emergency medical services (EMS).*
EMT. See *Emergency medical technician (EMT).*
EMTALA. See *Emergency Medical Treatment and Active Labor Act (EMTALA).*
EMT-B. See *Emergency medical technician-basic (EMT-B) curriculum.*
Encephalitis, Japanese. See *Japanese B encephalitis immunization.*
neuroimaging, 520
tickborne, immunization against, 61
Encephalopathy, anoxic, as poisoning complication, 436
Endocrinopathy, in potential organ donor patient, 720
Endometrioma, sonographic findings, 819
Endotracheal intubation, in prehospital setting, 284–285
confirming correct placement of, 286–287
beer in stomach and, 348–349
disposable calorimetric end-tidal carbon dioxide detector, 287
syringe aspiration technique, 287
ultrasound guiding, 905–907
Endovaginal sonography (EVS), female pelvis, 789
normal anatomy, 793–795
transabdominal ultrasonography versus, 790–792
Enhancement, ultrasound images, 955, 959–960
ENT. See *Ear, nose, and throat (ENT).*
Environmental emergencies, **451–476**
barotrauma, 457–460
carbon monoxide exposure, 465–468
decompression sickness, 455–457
high-altitude sickness, 451–455
hypothermia, 469–473
near drowning, 461–464
venomous snakebites (U.S.), 473–476
Environmental risks, to wilderness travel, altitude, 20–23, 206
air travel, 23
cardiovascular effects, 20–21
pulmonary effects, 21–23
barotrauma, 23–25, 226–229
cold, 18–19
heat, 19–20
infectious diseases, 25–26
malaria, 25–26
traveler's diarrhea, 25
vaccines, 26
SCUBA, 23–25, 226–229
Environmental somatization syndrome, 434
Epidemic hysteria, 434
Epiglottitis, in adult, and sore throat, case scenario, 335
Epinephrine, for pediatric upper airway infections, 383
racemic, for croup, hospitalization and, 330–331
Epistaxis, management advice, 336–337
Epstein-Barr virus disease, weakness associated with, 618
Ergotamine, for migraine, 407
for pediatric migraine, 631
Escherichia coli, causing traveler's diarrhea, 6, 182
Esophageal intubation , inadvertent or unrecognized, in prehospital setting, 286–287
Esophageal-tracheal double lumen airway, 284
Esotropia, 653, 663
Estradiol drips, for chest pain, 349
Ethnicity, undertreatment of pain and, 400
Evacuation, injured marine crew members, 246–247
insurance coverage, 4
Excitatory amino acid antagonists, in CNS resuscitation, 529–533
AMPA/KA receptor modulation, 532
glutamate release inhibition, 532–533
NMDA receptor, 530–532. See also *N-methyl-*D- aspartate (NMDA) receptor.
Excyclotorsion, 650
Exercise, at high altitude, 195
in pregnancy, 105–107
Exercise testing, for coronary artery disease patients, 31
Exophoria, 663
Exotropia, 663
Extended stay, immunization recommendations, 62–65
External chest compression, in asthma, 317
External ear barotrauma, 458
Extorsion, 650
Extracorporeal rewarming, hypothermic patient, 471
Extraocular muscles, 651–652, 653
actions and innervation, 652
diplopia and, 655–656
disorders, 655
Extrauterine empty gestational sac, 804

Extremities, pearls, radial head subluxation, 371–372
recognizing easily missed fractures and dislocations, lower extremities, 366, 369–371
upper extremities, 366, 367, 368
scapular manipulation of anterior shoulder dislocations, 365–366
tourniquet examination, 371
pitfalls, compartment syndromes, 373
fractures of nonaccidental trauma, 372
hip pain in children, 374–375
and spine, **365–377**
updates, acute knee injuries, decision rules for, 375–376
Ottawa ankle rules, 375
penetrating injuries, ultrasound evaluation of, 377
rotator cuff injury, ultrasonography and, 376
Eye, and intraocular structures, 649–650, 651
and ocular motor system, 650–652, 653
Eye drops, antibiotic, 332–333
beta blockers, for glaucoma, chronic heart failure and, 348
Eyepatch, with antibiotic therapy, 333

Fab fragment antidotes, 445
Facial nerves, functional grading system for, 332
in neurologic examination, 497–498
Falls, pregnant traveler, 97
Far field, ultrasound beam, 954
Fasciotomy, contraindicated for venomous snakebites, 474–475
Fatigue, chronic fatigue syndrome, 618, 619
weakness versus, evaluation in neurologic examination, 493, 493
Feet, bare, dangers of, 7–8
Fetal demise, ultrasound evaluation, 812–813
Fetal heart motion, sonographic detection, 799
Fetal hemoglobin, falsely elevated COHb levels and, 467
Fetal monitoring, following trauma to pregnant traveler, 97–98
Fetal pole, sonographic measurement, 798–799
Fetomaternal hemorrhage, 395–396
Fever, in traveler, differential diagnosis, 127
vaccine reaction, 263–264
Fibromyalgia syndrome, weakness associated with, 618, 619
classification criteria, 620
Fibrositis, weakness associated with, 618
Finger-to-nose testing, 502
First aid kit, commercial airline, 253
First trimester, vaginal bleeding in, ultrasound evaluation, 800–813
ectopic pregnancy. See *Ectopic pregnancy.*
incomplete abortion, 811–812
miscarriage, 811
threatened abortion, 811
viable versus nonviable fetus, 812–813
Fishing, medical emergencies and. See *Maritime medicine.*
Fish poisoning, **157–175,** 162–163. See also *Food poisoning.*
ciguatera, 165–168
scromboid, 161, 164
tetrodoxic or pufferfish, 168–170
weakness induced by, 617
Flatus, at high altitude, 206
Fluid collections, drainage, ultrasound guidance for, 908–909, 910
ultrasound characteristics, 922–923
Fluid percussion brain injury, 528–529
Fluid resuscitation, of trauma patients in prehospital setting, 296–297
Fluids, jet lag and, 12
in management of traveler's diarrhea, 185
Flumazenil, as antidote to benzodiazepine poisoning, 443–444
Focal neurologic deficits, at high altitude, 204
Focal zone, ultrasound beam, 954
Follicular cysts, sonographic findings, 819
Food, poisonings: food, fish, shellfish, **157–175.** See also *Food poisoning.*
precautions concerning, infants and children, 75–76
international travel and, 6
pregnant traveler, 108
Food poisoning, bacterial toxins involved, 158–159
botulism, 160–161
ciguatera, 165–168
diagnosis, 158, 159–160
incidence, 157–158
plant toxicity and, 173–174
scombroid, 161, 164
shellfish poisoning, 170–172
tetrodoxic or pufferfish poisoning, 168–170
treatment, 159–160
Foot, normal bony alignment rules, 369, 370
Foreign body(ies), esophageal, in pediatric patient, 386
in eye, antibiotic therapy for, 333
nasal, removal of, 328–329
and new onset wheezing in pediatric patients, 319–320

Foreign body(ies) *(Continued)*
removal during wound inspection, 421
ultrasound detection of, and procedure guidance, **895–910**
echo patterns, 897–901
localization and retrieval, 901–902
material and detection techniques, 895–897
potential applications in emergency department, 902, 904–910
safety issues, 902, 903–905
Fractures, cervical spine, 373–374
failure to recognize, lower extremities, 366, 369
upper extremities, in children, 366, 368
of nonaccidental trauma, 370, 372
diaphyseal, 372
metaphyseal, 372
Freedox. See *Tirilizad mesylate.*
Free radical scavengers, in CNS resuscitation, 533–534
Frequency, for ultrasound, commonly used, 937
and image resolution, 937, 949
sound waves, 924, 925
techniques, 971
Frequency shift, Doppler ultrasound, 859
Furazolidone, for diarrhea treatment in pregnant traveler, 109
Fusion (ocular), 663
failure, 657–658

GABHS. See *Group A beta-hemolytic strep (GABHS).*
Gain, ultrasound imaging, 942–944
Gain artifact, ultrasound images, 965, 967, 968
Gallbladder, sonographic evaluation, acoustic window for visualization, 826
normal appearance, 826–828
pathology, 828–834
polyps, 833–834
Gallstones, ultrasound images, 953, 955–959
with shadow effect, 899
sonographic patterns, 828–830, 835
Gangrenous cholecystitis, ultrasound findings in, gangrenous form, 831
Gases, behavior of, diving and, 225–226
in-flight atmosphere and, 253
Gastric irrigation, hypothermic patient, 471
Gastrointestinal disorders, air travel and, 257–258
maritime medicine and, 243–244
wilderness travel and, 35
Gastrointestinal injury, pediatric, foreign bodies causing, 386
Gastrointestinal symptoms, as travel vaccine reaction, 262–263
Gaze, conjugate, 654
GCS. See *Glasgow Coma Scale (GCS).*
Gender, undertreatment of pain and, 400
Genetic resistance, to malarial infection, 120
Gentamicin, for ophthalmologic problems, 333
Germicidal soap, 6
Gestation, normal versus abnormal prediction, 390
Gestational sac, extrauterine empty, 804
sonographic evaluation, 796–798
Giardia lamblia, causing traveler's diarrhea, 6, 183
Glasgow Coma Scale (GCS), in mental status evaluation, 494–495
near drowning victim and, prognosis, 463
withholding/withdrawing treatment, 464
in traumatic brain injury, classifying, 565, 567–568, 593, 595
defining, 563–564
mild form, 567–564
severe form, 581
Glass foreign body, ultrasound images, 898
Global amnesia, transient, 646
Glossopharyngeal nerve (cranial nerve IX), in neurologic examination, 499
Glucocorticoids, in CNS resuscitation, 534
Glucose, in stroke pathophysiology, 554
Glycine, and NMDA receptor activation, 531–532
"Good Samaritan" laws, 254
Gradenigo's syndrome, 656, 663
Gray scale, ultrasound image, 939–940
Group A beta-hemolytic strep (GABHS), 334–335
Grunting respirations, in pediatric patient, 384
Guillain-Barré syndrome, 610, 613, 657
Gulf War syndrome, weakness associated with, 618
Gynecologic emergencies. See *Obstetric and gynecologic emergencies.*

HACE. See *High-altitude cerebral edema (HACE).*
Haemophilus influenzae, in community-acquired pneumonia, outpatient therapy for, 320–321
type b immunization, in childhood, 48–49, 78
Haemophilus influenzae (HIB) vaccine, 383
HAFE. See *High-altitude flatus expulsion (HAFE).*

Halofantrine, coronary artery disease patients and, 33
Halogens, for water purification, 184
Hand grasp test, in neurologic examination, 499
Handicapped patients, wilderness travel and, 35–36
HAPE. See *High-altitude pulmonary edema (HAPE).*
Hazardous materials incidents, and psychologic injury, 433–434
Headache(s), in children, 625–627. See also *Migraine, pediatric.*
classification system, 406–407
differential diagnosis, 626
evaluation in neurologic examination, 493
incidence, 406
types, 626–627
Head trauma, classification, 565
and treatment, 569
mild, **563–576.** See also *Mild head trauma.*
severe brain injury, **581–600.** See also *Severe traumatic brain injury.*
Health care insurance, limitations when traveling, 4
Health care issues, for travelers, 5–13
access, 5
bare feet, 7–8
clothing, 9
driving, 6
food precautions, 6
insect repellents, 7
jet lag, 10–12
motion sickness, 12–13
motor vehicles, 6
sun, exposure to, 9–10
swimming, 8–9
water precautions, 6
Health history, of traveler, 18
Hearing impaired, communication with elderly, 337–338
Heart. See *Cardiac entries.*
Heat stroke, in serotonin syndrome differential diagnosis, 672
Heat tolerance, travel and, 19–20
Heel-to-shin testing, 502
Hemangioma, hepatic, ultrasound findings, 836
Hematologic changes, at high altitude, 194
Hematuria, in renal colic diagnosis, 402–403
Hemiplegic migraine, 628
Hemodialysis, hypothermic patient, 471
Hemodynamic compromise, in poisoned patient, 430–433
Hemopericardium, echocardiographic diagnosis, 759–761
Hemorrhage. See also *Bleeding.*
brain involvement. See *Brain hemorrhage.*
fetomaternal, 395–396
sonographic appearance, in blunt abdominal trauma, intra-abdominal, 771, 773
subdiaphragmatic, 767, 771
subarachnoid, misdiagnosis, 409–410
Hemorrhagic infarcts, 515, 554–555
Hemosalpinx, ectopic pregnancy and, 805, 806
Hemostasis, compensatory, in pediatric patient, 381
Hemothorax, sonographic appearance, in blunt abdominal trauma, 767
Henry's law, 225–226
Heparin, for stroke patient, 559
Hepatic cyst, ultrasound-guided drainage, 909, 910
Hepatic hemangioma, ultrasound findings, 836
Hepatitis A immunization, 52, 53, 56–57
for elderly, 66
for immunocompromised person, 66
for infants and children, 66
in pregnancy, 66, 99
Hepatitis B immunization, adult traveler, 53
for elderly, 66
extended stay recommendations, 62
for immunocompromised person, 66
for infants and children, 66
in pregnancy, 66, 99
routine, in childhood, 50–51, 79
Hepatocellular carcinoma, ultrasound findings, 836
Heteronymous diplopia, 652, 653, 663
Heterophoria, 663
Heterotopic pregnancy, sonographic appearance, 808–809
HIB vaccine, 383
High altitude, acclimatization to. See *Acclimatization.*
defined, 191–192, 192–193
traveler recommendations, 208. See also *High-altitude medicine.*
High-altitude bronchitis, 205
High-altitude cerebral edema (HACE), 191, 201–202, 454
treatment, 198, 201
High-altitude flatus expulsion (HAFE), 206
High-altitude medicine, **191–214**
accelerated acclimatization, 212
controversies in, 212–214
High-altitude pharyngitis, 205
High-altitude pulmonary edema (HAPE), 191, 202, 453–454
pathophysiology, 453
predisposing factors, 202–203
prevention, 204
treatment, 198, 201, 203–204, 453–454
High-altitude retinopathy, 205

High-altitude sickness, 451–455
acute mountain sickness, 452–453
cerebral edema, 454
pulmonary edema, 453–454
High-altitude syncope, 205
Himalayan hack, 205
Hip pain, in children, 374–375
Hismanal, causing long QT arrhythmias, 349
HIV-infected patients, wilderness travel and, 35
Holter monitoring, in diagnosis of spells, 645
Homonymous diplopia, 652, 653, 663
Hoover's sign, 504
Hospital admission, of croup patients treated with racemic epinephrine, 330–331
orders for, 310
of stroke patient, 558
HPS. See *Hypertrophic pyloric stenosis (HPS).*
5-HT. See *5-Hydroxytryptophan (5-HT).*
Human chorionic gonadotropin (hCG), measurement in pregnancy, 390
Human immunodeficiency virus (HIV)-infected patents, wilderness travel and, 35
Humidity, air travel and, 253
Hydration, jet lag and, 12
in management of traveler's diarrhea, 185
Hydrocephalus, neuroimaging, 522–523
Hydronephrosis, ultrasound image, 880, 888–890
5-Hydroxytryptophan (5-HT), serotonin syndrome and, 666–667
Hyperbaric oxygen therapy, for carbon monoxide exposure, 466
guidelines, 466
for decompression sickness, 231–234, 456–457
adverse effects, 232, 234
US Navy protocol, 233
for mountain sickness, 212–213
AMS, 197–198
HAPE, 203
portable chambers. See *Portable hyperbaric chambers.*
Hypercapnia, permissive, 322
Hypercarbia, in asthma and COPD, 318–319
Hyperechoic structure, on ultrasound image, 940
diaphragm, 942
Hyperkalemia, as cause of weakness, 611
Hypermagnesemia, as cause of weakness, 611
Hypersensitivity reaction, to travel vaccine, 262
Hypertension, high altitude travel and, 209
in potential organ donor patient, 721
Hypertonic saline, for blood pressure resuscitation, in severe traumatic brain injury, 592
Hypertrophic pyloric stenosis (HPS), ultrasound findings, 840–842
Hyperventilation, in high-altitude mountain sickness, 452
for severe traumatic brain injury patient, 599–600
Hypoglossal nerve (cranial nerve XII), in neurologic examination, 499
Hypokalemia, as cause of weakness, 611
Hypophosphatemia, as cause of weakness, 610
Hypotension, in potential organ donor patient, 720
of unknown cause, echocardiographic diagnosis, 756–757, 758
Hypothermia, 469–473
CNS resuscitation and, 534–535
management issues, 472–473
in potential organ donor patient, 720
resuscitation guidelines, 470
rewarming techniques, 470–472
in severe traumatic brain injury patient, 600
in special populations, 472
Hypovolemia, echocardiographic diagnosis, 756
recognizing in pediatric patient, 381

IAMAT. See *International Association of Medical Assistance to Travelers (IAMAT).*
Ibuprofen, for pediatric migraine, 631
Ice, contraindicated for venomous snakebites, 474–475
ICP. See *Intracranial pressure (ICP).*
IGF–1. See *Insulin-like growth factor–1 (IGF–1).*
IHS. See *International Headache Society (IHS).*
Illness, travel-related. See also *individual diseases.*
at altitude. See *Altitude sickness.*
epidemiology, 2
food causing. See *Food poisoning.*
in infants and children, 89–90
maritime. See *Maritime medicine.*
Immune suppression, at altitude, 206
Immunity, to malarial infection, 120
Immunizations. See also *Vaccines.*
maritime medicine and, 243
routine (childhood and adolescent), 44–46, 48–52
accelerated schedule, 47

and travel, **43–69**
adverse reactions, 2, 261–264. See also *Travel vaccine reactions.*
for children, 65, 66
country requirements, 59–61
for elderly, 66, 67
extended stay recommendations, 62–65
future vaccine possibilities, 67–68
geographic considerations, 61–62
for immunocompromised person, 66, 67
indications, 44, 45
in pregnancy, 65–67
risks and benefits, 44
routine requirements, 52–59
travel medicine consult and, 43–44
Immunocompromised traveler, immunizations for, 66, 67
wilderness travel and, 35
Immunoglobulin vaccine, 53
for elderly, 66
for immunocompromised person, 66
for infants and children, 66, 81–82
in pregnancy, 66
special groups recommendations, 66
Imodium, for traveler's diarrhea, 185, 187
Impedance plethysmography (IPG), in lower extremity deep vein thrombosis evaluation, 861–862
Incomplete abortion, ultrasound evaluation, 811–812
Infants and children, travel with, **71–90**
diseases acquired during, 86–89
malaria, 86–88
traveler's diarrhea, 88–89
immunizations, routine, 44–52, 78–80
travel-related, 65, 66, 77–78
post-trip follow-up, 89–90
pre-trip planning, 71–73
sample medical kit, 73
special considerations, eating and drinking, 75–76
recreation, 76–77
during travel, 74
wilderness travel, 28–30
Infarcts, bland, 515
hemorrhagic, 515, 554–555
lacunar, 516–517, 553
Infection(s). See also *specific infections.*
acute renal, formal ultrasound evaluation, 881–882
at altitude, 206
as CNS dysfunction cause, neuroimaging, 518–522, 523, 524
in AIDS patient, 520–521, 523–525
brain abscess, 518, 519
empyema, 518–519
encephalitis, 520
meningitis, 518
aseptic, 519
chronic, 520
headaches associated with, 626
maritime medicine and, 243
nonrespiratory, and grunting respiration in pediatric patient, 384
traveler and, 25–26
Inferior vena cava (IVC), in ultrasound examination of aorta, 852–853, 854
Infiltration, anesthetic, in wound care, 419
Inflammation, diplopia caused by, 658
headaches associated with, 626
inhibitors of, CNS resuscitation and, 535–536
In-flight deaths, causes, 252
Influenza immunization, 51, 53
for elderly, 66
Haemophilus influenzae (HIB) vaccine, 383
for immunocompromised person, 66
for infants and children, 66, 82
in pregnancy, 66
Infrared tympanic thermometers (ITTs), 304–305
Inner ear barotrauma, 457
Insect bites, weakness induced by, 617
Insect repellents, for travelers, 7
Insomnia, with or without acute mountain sickness, treatment, 200–201
Inspection, of wounds, 420–421
Instrumentation, for ultrasound, 930–937
Insulin-like growth factor–1 (IGF–1), in CNS resuscitation, 537–538
Insurance, and travel. See *Travel insurance.*
Intercostal approach, ultrasound scanning technique, 972
Interleukins, in CNS resuscitation, 535–536
International Association of Medical Assistance to Travelers (IAMAT), 5, 95
International Headache Society (IHS), diagnostic criteria for migraine in children under 15 years, 627–628
International travel. See *Travel; Travelers; Travel medicine.*
Internuclear ophthalmoplegia, 657
Interstitial ectopic pregnancy, sonographic appearance, 808
Intestinal nematodes, in post-travel patients, 275–277
Intracerebral hemorrhage, neuroimaging, 512–514
Intracranial injuries, delayed, in mild head trauma, 573
Intracranial neoplasms, neuroimaging, 523–524
Intracranial pressure (ICP), in severe traumatic brain injury, 585–588
management techniques, 598–600
Intradermal sutures, for wound closure, 422
Intraocular structures, diplopia and, 649–650

Intraosseous infusion-induced compartment syndromes, 373
Intraperitoneal fluid, in blunt abdominal trauma, sonographic appearance, 766–767
  minimal detectable quantity, 773
  scoring scale, 774
Intrauterine pregnancy, definite, 799, 800
  and ectopic pregnancy combined, 808–809
  no definite, 800, 803, 809
  probable abnormal, 800, 801
  in rule-out ectopic protocol, 800, 802
Intravenous fluids, very hot, hypothermic patient, 471–472
Intravenous pyelography (IVP), in nephrolithiasis diagnosis, 403
  for renal colic diagnosis, 879, 881
Intraventricular hemorrhage, neuroimaging, 514, 515
Intubation, endotracheal. See *Endotracheal intubation.*
Intussusception, ultrasound findings, 842–843
Iodine water purification tablets, for diarrhea prevention, 184
  use by pregnant traveler, 105
Ionophore blockade, NMDA receptor-associated, 530–531
Ipecac, syrup of, 296
IPG. See *Impedance plethysmography (IPG).*
Ipratropium bromide, for reactive airway disease, 315–316
Iritis, diagnosis, 333–334
Irrigation, of wound, 419–420
Ischemic cerebrovascular disease, classification, 514
  neuroimaging, 514–517
    ischemic strokes, 515–516
    lacunar infarcts, 516–517
Ischemic heart disease, echocardiographic diagnosis, 757–758
Ischemic strokes, neuroimaging, 514–516
Ischemic stroke syndromes, 553
Isolated calf-vein deep venous thrombosis, diagnostic ultrasound for, 868
Isometheptene, for pediatric migraine, 631
Isospora belli, causing traveler's diarrhea, 183
ITTs. See *Infrared tympanic thermometers (ITTs).*
IVP. See *Intravenous pyelography (IVP).*

Jamaican vomiting sickness, 174
Japanese B encephalitis immunization, 53, 62–63
  for elderly, 66
  for immunocompromised person, 66
  for infants and children, 66, 82–83
  in pregnancy, 66, 99
Jehovah's Witnesses, 312
Jet lag, 10–12
  management, 11–12
Johnson v. University of Chicago, 309
Joints, decompression sickness affecting, 456

Kernig's sign, 501
Ketamine, and NMDA receptor antagonism, 530–531
  pediatric use, 411–412
Ketorolac, for tension-type headache, 408
  in treatment of renal colic, 403–405
Khumbu cough, 205
Kidney. See also *Renal entries.*
  KUB radiography, in nephrolithiasis diagnosis, 403
  ultrasound image, 941–942. See also *Renal ultrasound.*
Kleinhauer-Betke test, in placental separation diagnosis, 394–395
Knee injuries, acute, decision rules for, 375–376
Kynurenate (KYNA), and NMDA receptor activation, 531–532

Labor, patient in, transfer rules, 287, 309–310
Laboratory toxidromes, 428
Lactose intolerance, in post-travel patients, 268
Lacunar infarcts, 553
  neuroimaging, 516–517
Lambert-Eaton myasthenic syndrome, 615
Lamotrigine, and glutamate release, 532
Landry-Guillain-Barré syndrome, 610, 613
Laparotomy, ultrasonography versus, in blunt abdominal trauma evaluation, 774–781
Larynx, tumors of, 329–330
Lateral cystic shadowing, ultrasound images, 924, 925
Lateral resolution, ultrasound images, 944, 947
  beam width on, 948
LEDVT. See *Lower extremity deep vein thrombosis (LEDVT).*
Left bundle branch block, myocardial infarction with, 344–345
Left parasternal windows, in 2-D echocardiography, 746–749
  normal view, 754
Legg-Perthes disease, in pediatric patient, 374

Leishmaniasis, cutaneous, in post-travel patients, 279
Leukoplakia, oropharyngeal, 329
Lidocaine, buffered versus unbuffered, for wound care, 417–418
Lights, use by emergency medical services vehicle, 291–293
Light touch, in sensory examination, 500
Limp, etiologic considerations in child, 374–375
Linear array scanner, ultrasound probe, 931–934
  curved, 934, 936
  focal length and direction of beam, 933
  image shape, 934
Linear electronic scanner, ultrasound probe, 931–934
Lipid peroxidation, CNS resuscitation and, 533–534
Lisfranc fracture dislocations, failure to recognize, 369–371
Liver, ultrasound findings, 826–827, 834–836, 942
  cirrhosis, 943
  pathology, 836
Loa loa infection, in post-travel patients, 275, 277
Longitudinal scan, ultrasound, 738–739
  kidney, 886, 887
Loperamide, for traveler's diarrhea, 185, 187
Lower extremity, ultrasound examination, 866–867
Lower extremity deep vein thrombosis (LEDVT), 860
  evaluation, 323–324
    by contrast venography, 860–861
    by impedance plethysmography, 861–862
    by magnetic resonance imaging, 867
    by radionuclide scanning, 867–868
    by ultrasonography, 862–865
Lumbar puncture, in diagnosis of spells, 644
  in pediatric meningitis evaluation, 384–385
  update on, 304
Lung disease, high altitude travel and, 208–209
Lung fluke infection, in post-travel patients, 273–274
Lungs, decompression sickness affecting, 456
Lyme disease, weakness associated with, 618

Magnesium, NMDA receptor modulation by, 531
  weakness induced by, 611, 616–617
Magnetic resonance (MR) angiography, in abdominal aortic aneurysm diagnosis, 851
  in stroke, 557
Magnetic resonance (MR) imaging, in central vein thrombosis evaluation, 870
  for cholelithiasis detection, ultrasound versus, 825
  in diplopia evaluation, 662
  for foreign body detection, 896
  in lower extremity deep vein thrombosis, 867
  in mild head injury, 574
  in nontraumatic CNS dysfunction, 509–510
    contrast media for, 510–511
  in pediatric migraine diagnosis, 630
  in vascular dissection evaluation, 692
Malaria, **113–150**
  acquired while traveling, clinical presentation, 122
    in infants and children, 86–88
    onset latency, 265
    in pregnant traveler, 28, 99–101
  clinical manifestations, 121, 122
    acute renal failure, 125
    acute respiratory distress syndrome, 125
    anemia, 126
    blackwater fever, 125
    bleeding, 126
    cerebral form, 124–125
    in emergency department, 264
    hypoglycemia, 125–126
    jaundice, 126
    lactic acidosis, 125–126
    severe or complicated, 123–124
    uncomplicated, 121, 123
  complications, 143
  diagnosis, 128–132
    differential diagnosis, 126–128
    difficulties, 114
    in emergency department, 264–265
  genetic resistance, immunity, and premunition, 120–121
  global distribution, 113, 114
  incidence, 113
  parasite life cycle, 115–119
  parasitic resistance to antimalarial drugs, 113–114
  pathogenesis, 119–120
  prevention, 25–26, 143–149
    chemoprophylaxis, 145–148
    in pregnant traveler, 100–101, 104–105
    protection against mosquitoes, 143, 145
    vaccine, 148–149
  treatment, 101

Malaria *(Continued)*
antimalarial drugs, 132–138
children, 141–142
chloroquine-resistant *P. falciparum* and *P. vivax*, 139–140
chloroquine-sensitive malaria, 138–139
in emergency department, 265–267
exchange transfusion, 142
life-threatening malaria, 140–141
monitoring, 142
multidrug-resistant *P. falciparum*, 140
in pregnancy, 104–105, 141
severe malaria complications, 143
triage, 132–133
uncomplicated malaria, 138
Malaria parasite, life cycle, asexual liver and blood stages, 117–119
parasite, 115–116
sexual cycle in mosquito, 119
vector, 116–117
Malaria vaccine, 148–149
Malignancies, of head and neck, identifying, 329–330
Malingering, suspected, neurologic examination in, 503–505
weakness associated with, 619
Malpractice, admission orders and, 310
Mannitol, for severe traumatic brain injury patient, 591, 599
Mannkopf's sign, 504
MAOIs. See *Monoamine oxidase inhibitors (MAOIs).*
Maritime medicine, **241–248**
defined, 241
historical aspects, 241–242
scope of medical occurrences, 241–245
cardiovascular problems, 244
dehydration, 244
frequency, 242
gastrointestinal disorders, 243–244
immunizations and, 243
infectious diseases, 243
injury and illness patterns, 244–245
medications prescribed, 242–243
musculoskeletal problems, 242
recreational boating injuries, 245
telemedicine service role in, 245–248. See also *Telemedicine services.*
Masses, pelvic, ectopic pregnancy and, 805, 807, 808
sonographic findings, 818
endometriomas, 819
neoplasms, 820
simple cysts, 819
tubo-ovarian abscesses, 817, 818
renal, formal ultrasound evaluation, 882
Mass psychogenic illness, 434
MDI-spacer therapy, for bronchospastic patient, 322
Measles immunization, adult, 54
for elderly, 66
for immunocompromised person, 66
for infants and children, 66
pregnancy and, 66, 98
routine, in childhood, 49, 79
Mechanical sector scanner, ultrasound probe, 931, 932
image shape, 935
linear array scanner versus, 934
Medial longitudinal fasciculus (MLF), 654, 657, 659
Medical care, in foreign countries, pregnant traveler and, 95–96
Medical emergencies, in commercial air travel, **251–259.** See also *Commercial air travel.*
diving-related, **223–237**
maritime. See *Maritime medicine.*
obstetric and gynecologic. See *Obstetric and gynecologic emergencies.*
travel-related, timing related to, 2
Medical equipment, for airline personnel, 253–254
Medical evacuation, insurance covering, 4
Medical kits, for children and infants, 73
on commercial airlines, 253–254
diarrhea travel kit, 184
medication recommendations, 3
Medical oversight, lacking in emergency medical services education and practice, 293–295
Medical training, airline personnel, 253–254
telemedicine services, 247
Medications, adverse reactions to, in elderly, 437–438, 439
high incidence agents, 439
recognizing, 437–438
recommendations, 438
Medicolegal risk, admission orders and, 310
Medullary syndromes, clinical features, 687
Mefloquine therapy, contraindications, 25, 26
coronary artery disease patients and, 33
for malaria prevention, 145, 147, 148
in pregnant traveler, 100
Melatonin, jet lag and, 12
Melioidosis, in post-travel patients, 274
Meningitis, neuroimaging in, 518
aseptic form, 519
chronic form, 520
in pediatric patient, 384–385
Meningococcus immunization, 54
country requirements, 59
for elderly, 66
geographic considerations, 61
for immunocompromised person, 66
for infants and children, 66, 83

in pregnancy, 66, 99
special groups recommendations, 66
Mental stability, altitude and, 206
wilderness travel and, 37
Mental status, altered, patient history of, 492
in differentiation of spells, 641
in neurologic examination, 494–495, 496
Meperidine, for tension-type headache, 408
Merchant marine, medical services for. See *Telemedicine services.*
Metabolic acidosis, 428
in poisoned patient, adrenergic response to, 428–429
Metal foreign body, ultrasound images, 898
Metaphyseal long-bone fractures, nonaccidental, in children, 370, 372
Metastasis, spinal, 708–709
Methotrexate, in treatment of ectopic pregnancy, 393–394
N-methyl-D- aspartate (NMDA) receptor, competitive antagonism, 530
glycine site modulation and, 531–532
ifenprodil-like, 532
ionophore blockade associated with, 530–531
magnesium modulating, 531
in stroke pathophysiology, 554
Methylprednisolone, in CNS resuscitation, 534
4-Methylpyrazole (4-MP), as antidote to methanol and ethylene glycol intoxication, 444–445
Metoclopramide, for migraine, 408–409
for pediatric migraine, 631
serotonin syndrome and, 670
Midazolam, for pediatric patient, 385–386
Middle ear squeeze, 457
Migraine, 646
compazine for, 408–409
ergotamine for, 407
ketorolac versus meperidine for, 408
pediatric, **625–634**
complicated, 628
criteria, 627–628
diagnostic studies, 629–630
differential diagnosis, 625–627
features, 628
pathophysiology, 629
physical examination in, 629
prognosis, 634
treatment, 630–634
reglan for, 408–409
serotonin and, 629
subarachnoid hemorrhage misdiagnosed as, 409–410
sumatriptan for, 407
adverse effects, 407–408
traumatic head injury inducing, 675
Mild head trauma, **563–576**
classification of head injury, 565
and treatment, 569
defined, 563–564
delayed intracranial injuries, 573
epidemiology, 564
evaluation and management, 565–573
MR imaging in, 574
pathophysiology, 564
postconcussion syndrome in, 574–575
sports injuries and concussions, 575, 576
Mini-mental status examination, 495, 496
Minors, "mature," 311
and refusal of care, 311–312
treatment, 310–311
Mirror effect artifact, ultrasound images, 965, 966–967, 968
Miscarriage, ultrasound evaluation, 811
MK801, and NMDA receptor antagonism, 530–531
MLF. See *Medial longitudinal fasciculus (MLF).*
M mode echocardiography, 940
Monitoring, in severe traumatic brain injury, 595–596
blood pressure, 592
Monoamine oxidase inhibitors (MAOIs), overdose, 442
serotonin syndrome and, 665–666, 667–668
Monocular diplopia, 658–659
defined, 663
transient unilateral, 659, 660
Monro-Kellie doctrine, 585
Morison's pouch, sonographic appearance, in blunt abdominal trauma, 767, 768–770, 773
Morphine, pediatric use, 411
for renal colic pain management, 406
Mosquito bites, protection against, 7, 100, 143, 145
Motion sickness, preventive strategies, 13
treatment for pregnant traveler, 103–104
Motor function, in neurologic examination, 499–500
Motor vehicles, accident involving pregnant traveler, 96–97
international travel and, 6
"Mountaineer's rule," 199
Mountain sickness, acute. See *Acute mountain sickness (AMS).*
chronic, 207
subacute, 207
4-MP. See *4-Methylpyrazole (4-MP).*
MR imaging. See *Magnetic resonance (MR) angiography; Magnetic resonance (MR) imaging.*
Mumps immunization, pregnancy and, 98
routine, in childhood, 79
Munchausen's syndrome, 618

Muscle contraction headaches, 626
Muscles, diplopia and, 661
extraocular, 651–652
ocular motor system, 650–652, 653
Muscle structures, female pelvis, transabdominal ultrasonography examination, 793
Muscle weakness. See *Weakness.*
Musculoskeletal disorders, maritime medicine and, 242, 245
wilderness travel and, 35
Mushroom poisoning, 173
Myasthenia gravis, weakness in, 611, 614–615
Mycoplasma pneumoniae, bedside cold agglutinin test for, 316–317
in community-acquired pneumonia, outpatient therapy for, 320–321
Myiasis, cutaneous, in post-travel patients, 278
Myocardial infarction, acute. See *Acute myocardial infarction (AMI).*
with bundle branch block, diagnosis, 344–345
diagnosis of, pacemaker and, 342
Myofascial pain syndrome, 618
Myopathy, weakness associated with, 608. See also *Weakness.*

NAC. See *N-acetylcysteine (NAC) therapy.*
NAEMSP. See *National Association of Emergency Medical Service Physicians (NAEMSP).*
Nalmefene, as antidote to opiate toxicity, 444
Naloxone, as antidote to opiate toxicity, 444
Narcotics, for abdominal pain, 405–406
Nasal cavity, visualization techniques, 336, 337
Nasopharynx, malignancies of, 329
National Acute Spinal Cord Injury Study (NASCIS), 534
National Association of Emergency Medical Service Physicians (NAEMSP), emergency vehicle use recommendations, 293
National Highway Traffic Safety Administration (NHTSA), new EMT-B curriculum and, 295–296
Natural immunity, to malarial infection, 120
Near drowning, 461–465
epidemiology, 461
treatment guidelines, 461–469
ACLS guidelines, 461–462
outpatient management, 462
surfactant treatment, 462
survival prognosis, 463
withholding/withdrawing treatment, 463–464
Near field, ultrasound beam, 954
Neck masses, children versus adults, 330
evaluation difficulties, 330
symptoms associated with, 330
Nefazodone overdose, 443
Nematodes, intestinal, in post-travel patients, 275–277
Neomycin, for ophthalmologic problems, 333
for suture line aftercare, 424
Neoplasms, intracranial, neuroimaging, 523–524
ultrasound findings, biliary tract, 833–834
female pelvis, 820
liver, 836
Nephrolithiasis, diagnosis, 402–403
imaging modalities, 403
Nephropathy, from contrast agents, 510
Nerve growth factor (NGF), in CNS resuscitation, 536–537
Nerves, cranial. See *Cranial nerves.*
spinal cord, 700
Neural mismatch hypothesis, motion sickness and, 12
Neuroimaging, **507–525**
in nontraumatic CNS dysfunction, 507–525
causes, 511–525
brain hemorrhage, 511–514
diffuse cerebral edema, 517–518
hydrocephalus, 522–523
infection, 518–522, 523, 524
intracranial neoplasms, 523–524
ischemic cerebrovascular disease, 514–517
contrast imaging in, 510–511
differential diagnosis, 507, 508
methods, 507–510
Neuroleptic malignant syndrome (NMS), in serotonin syndrome differential diagnosis, 671–672
Neuroleptics, toxicology of, 442–443
Neurologic decompression sickness, 456
Neurologic deficits, as DCS manifestation, 230–231
at high altitude, 204
wilderness travel and, 36
Neurologic examination, **491–506**
in comatose patient, 502–503
coordination and balance in, 502
cranial nerves in, 495–499
directed versus complete, 492
documenting, 505–506
history in, 492–494
mental status in, 494–495, 496
motor testing in, 499–500

reasons for, 491
reflexes in, 501
sensory testing in, 500–501
in spinal cord syndromes, 701–703
in suspected conversion disorder or malingering, 503–505
Neurologic shellfish poisoning (NSP), 163, 171–172
Neuromuscular blockade (NMB), prehospital use, 298–299
contraindications, 298
EMT training for, 299
ideal agent, 299
for severe traumatic brain injury patient, 598
Neuromuscular disease, peripheral, approach to generalized weakness and, **605–622.** See also *Weakness.*
Neuropathies, peripheral, weakness associated with, 608
Neurotrophic factors, CNS resuscitation and, 536–538
NGF. See *Nerve growth factor (NGF).*
Nifedipine, for acute mountain sickness, 213
for HAPE, 201, 203
Nimodipine, in CNS resuscitation, 528
Nitrates, administration techniques in acute pulmonary edema, 349
Nitrous oxide, pediatric use, 411
NMB. See *Neuromuscular blockade (NMB).*
NMDA. See *N-methyl-*D- aspartate (NMDA) receptor.
NMS. See *Neuroleptic malignant syndrome (NMS).*
Nonaccidental trauma, fractures associated with, 370, 372
Noninvasive ventilatory support, 321
Nonrespiratory infections, and grunting respiration in pediatric patient, 384
Non-steroidal anti-inflammatory drugs (NSAIDs), pediatric use, 411
in treatment of renal colic, 403–405
Norwalk virus, causing traveler's diarrhea, 182
Nose, malignant lesions of, 329
Nosebleeds, management advice, 336–337
Noxious stimulus, 504–505
NSAIDs. See *Non-steroidal anti-inflammatory drugs (NSAIDs).*
NSP. See *Neurologic shellfish poisoning (NSP).*
Nuclear scintigraphy, for coronary artery disease diagnosis, 352–354
Numbness, evaluation in neurologic examination, 493, 493
Nursemaids' elbow, 371–372
Nylen-Barany maneuver, vestibular nerve and, 498
Obesity, wilderness travel and, 36
Oblique scanning, ultrasound examination of the aorta, 854–855, 856
Obstetric and gynecologic emergencies, indications, 96
pearls, ectopic pregnancy, criteria for further evaluation, 390–391
pitfalls, and updates, **389–396**
pitfalls, ectopic pregnancy, failure to screen for, 389
ruptured, indicator reliability of, 389–390
pelvic ultrasonography usefulness, 390
serum hCG in normal or abnormal gestation prediction, 390
updates, ectopic pregnancy, methotrexate treatment for, 393–394
serologic markers of, 391–392
traumatic placental separation, diagnosis of, 394–395
emergency department ultrasound, 392–393
fetomaternal hemorrhage, 395–396
Obstruction, ultrasound diagnosis, biliary system, 831
bowel, 839
hypertrophic pyloric stenosis, 840–842
Obstructive lung disease, ventilatory management in patients with, 318
Occult reflex, 501
OCG. See *Oral cholecystography (OCG).*
Ocular imbalance, headaches associated with, 627
Ocular motor system, cranial nerves in, 652–654
muscles in, 650–652, 653
supranuclear control, 654
Oculomotor nerve (cranial nerve III), diplopia and, 652–653, 656–657
in neurologic examination, 496–497
OEII chamber experiment, 20–21
Olfactory nerve (cranial nerve I), in neurologic examination, 495
Oligoanalgesia, 399–400
Onboard medical kits, commercial airlines, 253–254
Ondansetron, serotonin syndrome and, 670
ONO–3144, in CNS resuscitation, 533
Operational procedures, in-flight medical emergencies, 254–255
Operation Everest II (OEII) chamber experiment, 20–21
Ophthalmologic problems, therapeutic approaches, 332–333
Ophthalmoplegia, internuclear, 657
Ophthalmoplegic migraine, 628
Ophthalmoscopy, 660
Optic nerve (cranial nerve II), in neurologic examination, 496
Oral cholecystography (OCG), for cholelithiasis detection, ultrasound versus, 825–826

Oral rehydration solution (ORS), in management of traveler's diarrhea, 185
Organ donor, potential, care of, 720–721
Organophosphates, as cause of weakness, 611
Organ transplantation, following carbon monoxide exposure, 467
Oropharynx, malignancies of, 329
Osmolality, of contrast agents, 510
OTC medications. See *Over-the-counter (OTC) medications.*
Otitis media, 327–328
  standard treatment recommendations, 327–328
Ottawa ankle rules, 375
Outpatient therapy, for community-acquired pneumonia, 320–321
Ovaries, ultrasonography examination, endovaginal, 796
  torsion, 820–821
  transabdominal, 793
Overdose, in serotonin syndrome differential diagnosis, 672
Over-the-counter (OTC) medications, adverse reactions to, 437
  for travelers, 3
  use by elderly, 437
Oxford Scale, 500
Oximetry, in pediatric patient, 382–383
Oxygen delivery, effects of poisoning on, 431, 432
  therapeutic options to improve, 431, 433
Oxygen therapy, for hemodynamically compromised poisoned patient, 433
  for mountain sickness, 212
    AMS, 197
    HAPE, 203
  for severe traumatic brain injury, 588–590
  for stroke patient, 558

Pacemakers, and acute myocardial infarction diagnosis, 342
  and automated implanted defibrillator interaction. See *Automated implanted defibrillators.*
  tachycardia mediated by, 349–350
Pain, abdominal, in post-travel patient, 268
  achy limb, as DCS manifestation, 230
  management. See *Pain management.*
  pelvic, sonographic evaluation, ovarian torsion, 820–821
    pelvic inflammatory disease, 816–818
    pelvic masses, 818–820
    in pregnant trauma patient, 816
    vaginal bleeding associated with, in first trimester, 800–813
    in second and third trimesters, 813–815
  and swelling, as travel vaccine reaction, 263
Painful stimulus, in comatose patient, 503
  in sensory examination, 500
Pain management, pearls, migraine, compazine and reglan for, 408–409
    sumatriptan for, 407–408
  pitfalls, and updates for, **399–412**
  renal colic, ketorolac for, 403–405
    nephrolithiasis diagnosis and, 402–403
  tension-type headache, ketorolac and meperidine therapy compared, 408
  vaso-occlusive disease, patient controlled analgesia for, 401–402
  pitfalls, failure to recognize in children, 410–412
    migraine misdiagnosis, 409–410
    oligoanalgesia, 399–400
    renal colic misdiagnosis, 405
    sickle cell disease, inadequate treatment for, 401
    undertreatment, ethnicity and gender as risk factor for, 400
  updates, renal colic, narcotics for, 405–406
Papaveretum, for renal colic pain management, 406
Parallelism, carpal dislocations and, 365, 366
Paralytic shellfish poisoning (PSP), 163, 171–172
Paraovarian cysts, sonographic findings, 819
Parasites, bronchospasm induced by, in post-travel patients, 274–275
  in post-travel patients, 275–277
Parinaud's syndrome, 657, 663
Parks-Bielschowsky three-step test, 661
Parotid gland tumors, 329
Partial cord syndromes, 373–374
PASG. See *Pneumatic antishock garment (PASG).*
Passive rewarming, of hypothermic patient, 470
Patient, care refused by. See *Patient refusal.*
  transfer. See *Transfer of patients.*
  ultrasound examination, interaction with physician, 969–970
    moving prior to, 970
Patient-controlled analgesia (PCA), for vaso-occlusive crisis, 401–402
Patient dumping, prevention of, 309
Patient history, in neurologic examination, 492–494
Patient refusal, of care, dilemma for physician, 289–290

Jehovah's Witnesses, 312
minor patient and, 311–312
valid release for, 290
limited role of, 290–291
PCA. See *Patient-controlled analgesia (PCA).*
PCP. See *Phencyclidine (PCP).*
PEA. See *Pulseless electrical activity (PEA).*
Pearls, pitfalls, and updates, cardiology, **341–361**
clinical practice of emergency medicine, **303–312**
ear, nose, and throat, **327–338**
environmental emergencies, **451–476**
extremities and spine, **365–377**
obstetric and gynecologic emergencies, **389–396**
for pain management, **399–412**
pediatrics, **381–387**
prehospital care, **283–299**
respiratory care, **315–325**
in toxicology, **427–445**
wound care, **417–424**
Pediatric migraine, **625–634.** See also *Migraine, pediatric.*
Pediatric patient, abdominal ultrasound in, 840
blunt trauma to abdomen, 781–782
hypertrophic pyloric stenosis, 840–843
intussusception, 842–843
brain death in, 719
chloral hydrate use with, 386, 412
fractures of nonaccidental trauma in, 370, 372
headaches in, 625–627. See also *Migraine, pediatric.*
hip pain in, 374–375
hypothermia in, 472
infants and, travel with. See *Infants and children, travel with.*
missed supracondylar fractures in, 366, 368
new onset wheezing in, foreign body and, 319–320
pain in, failure to recognize and treat, 410–412
radial head subluxation in, 371–372
Pediatrics. See also *Pediatric patient.*
pearls, maximizing access, minimizing procedures, 382
pitfalls, and updates, **381–387**
recognizing stability, 381
rectal anticonvulsants, 385
sedation and analgesia, 385–386
ventilatory-perfusion mismatch, 383
pitfalls, aging bruises, 386–387
failure to recognize treat pain, 410–412
grunting respiration, 384
temperature measurement, 382
updates, esophageal foreign bodies, 386
evolving meningitis, 384–385
oximetry, 382–383
upper airway infections, 383
PEG. See *Polyethylene glycol (PEG).*
Pelvic inflammatory disease, sonographic findings, 816–818
Pelvic masses, sonographic findings, 818–820
Pelvic ultrasound, **789–821**
anatomy (normal pelvis), by endovaginal technique, 793–796
by transabdominal technique, 792–793
in early normal intrauterine pregnancy, confirming, 390
fetal pole, 798–799
gestational sac, 796–798
yolk sac, 798
in Emergency department, 392–393
endovaginal technique, 793–796
in first-trimester vaginal bleeding, ectopic pregnancy, 800–811
fetal demise, 812–813
incomplete abortion, 811–812
threatened abortion/miscarriage, 811
ovarian torsion, 820–821
pelvic inflammatory disease, 816–818
pelvic masses, 818–820
in pregnant trauma patient, 815
for pregnant traveler, following trauma, 98
prior to travel, 95
in second- and third-trimester vaginal bleeding, placental abruption, 814–815
placenta previa, 813–814
transabdominal, 390, 392
versus endovaginal, 790–792
technique, 792–793
transvaginal, 392
Penetrating trauma, to chest, transesophageal echocardiography in, 360–361
to extremities, ultrasonography evaluation, 377
to head and neck, carotid and vertebral artery dissections, 690–691
to heart, echocardiographic diagnosis, 759–761
Penicillin, for streptococcal pharyngitis, 334–335
Penn Travel Medicine (University of Pennsylvania Health System), insurance recommendations, 5
pre-trip planning suggestions, 3–4
Penumbra, in stroke pathophysiology, 553–554
Pepto Bismol. See *Bismuth subsalicylate.*
Percutaneous transluminal coronary angioplasty (PTCA), thrombolytic therapy versus, for acute myocardial infarction, 354–357

Pericarditis, colchicine relieving symptoms in, 348
Peripheral neuromuscular disease, approach to generalized weakness and, **605–622.** See also *Weakness.*
versus CNS disease, 605–606
Peripheral neuropathies, weakness associated with, 608
Peritoneal lavage, hypothermic patient, 471
Peritoneum, free fluid collection. See *Intraperitoneal fluid.*
Permethrin insecticide, 7, 100
Permissive hypercapnia (PHC), 322
Persistent postconcussive syndrome (PPCS), 574
Pertussis immunization, routine, in childhood, 48, 78
PET scans. See *Positron emission tomography (PET) scans.*
Pharmacologic intervention, jet lag and, 11–12
Pharyngitis, at high altitude, 205
PHC. See *Permissive hypercapnia (PHC).*
Phencyclidine (PCP), and NMDA receptor antagonism, 530–531
Photoallergic reactions, 10
Physical disabilities, wilderness travel and, 35–36
Physical examination, ultrasound scanning during, 971
"Physician certification for transfer," 289
Physician-patient interactions, 305–306
angry patient or family, 306–308
elderly patient, 337–338
positive behaviors, 306
during ultrasound examination, 969–970
Physician-physician interactions, 308–309
Physicians, renal ultrasound performed by, 884–891
role during in-flight emergencies, 255
Piezoelectric effect, 930–931
and image translation, 937–944
and transducer types, 931–937
Pill ingestion, ultrasound scanning, 904–905
Pinealoma, 657
Pinprick test, 702
Pit vipers, 473–474
envenomation by, antivenin for, 475–476
Placebo effect, migraine treatment in children, 630
Placental abruption, ultrasound evaluation, 814–815
in trauma patient, 816
Placental separation, diagnosis, 394–395
Placenta previa, ultrasound evaluation, 813–814
Plague, immunization against, 54, 63
in pregnant traveler, 99
Plantar-flexion reflex, 501
Plant toxicity, 173–174
ackee, 173–174
cassava, 174
Plasmodium, characteristics, 116
life cycle, 115–119
asexual liver and blood stages, 117–119
sexual cycle in mosquito, 119
vector, 116–117
Plasmodium falciparum, chloroquine-resistant, 139–140
complications from, 124
global drug resistance, 114
chemoprophylaxis recommendations, 144
lethality, 119–120
multidrug-resistant, 140
Plasmodium vivax, chloroquine-resistant, 139–140
Plastic foreign body, ultrasound image, 900
Platelet changes, at high altitude, 204
Pleural effusions, in chronic heart failure, 348
Pleural irrigation, hypothermic patient, 471
PMN. See *Polymorphonuclear (PMN) leukocytes.*
PMR. See *Polymyalgia rheumatica (PMR).*
Pneumatic antishock garment (PASG), limited use, 296
Pneumococcus immunization, 54
for elderly, 66
for immunocompromised person, 66
for infants and children, 51–52, 66, 83–84
in pregnancy, 66
Pneumomediastinum, in adult, and sore throat, case scenario, 335–336
Pneumonia, community-acquired, antibiotic choice for outpatient therapy, 320–321
Pneumonitis, aspiration type, as poisoning complication, 435
Poisoned patient, *N*-acetylcysteine therapy for, 441–442
antidepressants, 441–442
complications in, 435–436
anoxic encephalopathy, 436
aspiration pneumonitis, 435
compartment syndrome, 436
rhabdomyolysis, 436
evaluation of, laboratory role in, 427–430
hemodynamically compromised, approach to, 430–433
neuroleptics, 442
Poisoning. See *Food poisoning; Poisoned patient.*
Poliovirus immunization, adult, 54
for elderly, 66
for immunocompromised person, 66

for infants and children, 66
in pregnancy, 66, 99
routine, in childhood, 49–50, 79–80
Polyethylene glycol (PEG), in CNS resuscitation, 533–534
Polymorphonuclear (PMN) leukocytes, in CNS resuscitation, 535–536
Polymyalgia rheumatica (PMR), 618, 620
Polymyositis, weakness in, 611
Polyps, gallbladder, ultrasound findings, 833–834
Popliteal vein, in ultrasound examination of lower extremity, 866
Portable hyperbaric chambers, for mountain sickness, 212–213
AMS, 197–198
HAPE, 203
Position, patient, for ultrasound examination of the aorta, 854
transducer, for ultrasound evaluation of blunt abdominal trauma, 766
Positional arrhythmias, 348
Positive airway pressure, bilevel versus continuous, for acute pulmonary edema, 321
Positron emission tomography (PET) scans, in stroke syndromes, 553
Postconcussion syndrome, 574–575
Posterior tibial veins (PTV), in ultrasound examination of lower extremity, 866
Post-trip evaluation, 13–14
Posture, ultrasound scanning technique, 972
Potassium imbalance, weakness induced by, 611, 616–617
Povidone iodine, for wound irrigation, 420
PPCS. See *Persistent postconcussive syndrome (PPCS).*
PRCA. See *Percutaneous transluminal coronary angioplasty (PTCA).*
Predictive value, of diagnostic tests, 304
Pregnancy. See also *Pregnant traveler.*
ectopic. See *Ectopic pregnancy.*
hCG measurement in, 390
heterotopic, 808–809
high altitude travel and, 209–210
and travel, **93–109.** See also *Pregnant traveler.*
limitations, 93–94
ultrasonography confirming, 390
Pregnant traveler, airflight, 28, 107–108
drug use by, 101–105
for altitude sickness, 102
analgesics, 101
antidiarrheal and antiemetic, 101–102
antimalarial, 104
for infectious diseases, 104–105
insecticides, 105
iodine water purification tablets, 105
for motion sickness, coughs, and colds, 102–104
sleeping pills and tranquilizers, 102
eating and drinking considerations, 108
exercise and sports activity by, 105–107
at high altitude, 107
immunizations/vaccines for, 27–28, 65–67, 98–99
malarial infection in, 28, 99–101
medical care while abroad, 95–96
obstetric emergencies, 96
prenatal checkups, 94–95
pretravel checklist, 94
trauma to, 96–98
traveler's diarrhea in, 108–109
wilderness travel and, 26–28
Prehospital care, pearls, airway management for basic level provider, 283–285
confirming endotracheal tube placement, 286–287
early defibrillation and EMS system design, 285–286
pitfalls, and updates, **283–299**
pitfalls, COBRA transfers, 287–289
emergency vehicle lights and siren, use of, 291–293
lack of medical oversight, 293–295
patient refusals, 289–291
updates, cessation of resuscitation efforts, 297–298
fluid resuscitation of trauma patients, 296–297
neuromuscular blockers, use of, 298–299
new EMT-B curriculum, 295–296
Premunition, malarial infection, 120
Prenatal checkups, pregnant traveler and, 94–95
Pressure, atmospheric versus barometric, 192
diving and, 224
versus volume at depth, 225
Pressure equivalents, 224
Pressurization, airplane cabin, 210
in-flight physiology and pathophysiology and, 252–253
Pre-trip planning, 3–4
children and infants, 71–73
environmental risks and, 17–18. See also *Wilderness travel.*
pregnant traveler checklist, 94
Primaquine, malarial infection in pregnant traveler and, 100
Prioritization, of CT scanning, in severe traumatic brain injury patients, 596–598
Proarrhythmias, from antiarrhythmics, 349
Probe. See *Transducer.*
Procedures, foreign body ultrasound detection and guidance, **895–910**
echo patterns, 897–901

Procedures *(Continued)*
localization and retrieval, 901–902
material and detection techniques, 895–897
potential applications in emergency department, 902, 904–910
safety issues, 902, 903–905
minimizing in pediatric patient, 382
operational, in-flight medical emergencies, 254–255
surgical procedures, ultrasound use in, 742–743
Prochlorperazine, for acute mountain sickness, 197
for migraine, 408–409
pediatric patient, 631
Progesterone testing, in ectopic pregnancy diagnosis, 391–392
Proguanil, for malaria prevention, 145, 146–147
in pregnant traveler, 100
Projection (ocular), 663
Prolactin levels, in diagnosis of spells, 643
Promethazine, for pediatric migraine, 631
Prone scapular manipulation method, anterior shoulder dislocation, 365
Prophylaxis, for malaria prevention. See *Chemoprophylaxis.*
migraine treatment in children, 633
traveler's diarrhea, 187
Propranolol prophylaxis, for pediatric migraine, 633
Proprioception, in sensory examination, 500
Protozoal infestation, causing traveler's diarrhea, 183
Pruritus, as DCS manifestation, 230
Pseudobicornuate ectopic pregnancy, 809
Pseudogestational sac, ectopic pregnancy, 809, 810
Pseudo sludge, on ultrasound images, 961–962
PSP. See *Paralytic shellfish poisoning (PSP).*
Psychiatric disorders, altitude and, 206
weakness in, 618
wilderness travel and, 37
Psychogenic disorders, weakness associated with, 618–620
Psychogenic-induced symptoms, following hazardous materials incidents, 434
Psychologic injury, hazardous materials incidents and, 433–434
Psychosis, clinical characteristics, 494
evaluation in neurologic examination, 494–495
Psychotropic medicines, toxicology of, 442–443
PTV. See *Posterior tibial veins (PTV).*
Pufferfish poisoning, 162
Pulmonary barotrauma, 458–459
Pulmonary decompression sickness, 456
Pulmonary disorders, air travel and, 255–256
wilderness travel and, 34
Pulmonary edema, acute cardiogenic, bi-PAP versus CPAP for, 321–322
immediate nitrate administration techniques for, 349
at high altitude. See *High-altitude pulmonary edema (HAPE).*
Pulmonary embolism, during air travel, 258
suspected, evaluation of, 323–324
thrombolysis in, new indications for, 324–325
Pulmonary system, altitude and, 21–23
Pulse dampening, on axial resolution, ultrasound images, 951
Pulse duration, on axial resolution, ultrasound images, 950
Pulseless electrical activity (PEA), echocardiographic diagnosis, 755–756
Pulse oximetry, in carbon monoxide exposure, 467
in pediatric patient, 382–383
Pupillary reaction, in comatose patient, 503
in diplopia evaluation, 660
in neurologic examination, 496–497
Purification tablets, for water, use by pregnant traveler, 105
Pyloric stenosis, hypertrophic, ultrasound findings, 840–842
Pyrimethamine-sulfadoxine, for malaria prevention, 147, 148

QRS rhythms, regular wide, 345–347
VT/wide stable arrhythmias, 341–342
Quality assurance, in emergency ultrasound, 916–917
Quinidine, coronary artery disease patients and, 33
for malaria treatment, 267
Quinine, causing long QT arrhythmias, 349
for malaria treatment, 266, 267
Quinolones, pregnant traveler and, diarrhea treatment, 108

Rabies, immunization against, 55, 63–64
for elderly, 66
for immunocompromised person, 66
for infants and children, 66, 84
in pregnancy, 66, 99
Racemic epinephrine, for croup, hospitalization and, 330–331
Radial head subluxation, in children, 371–372

Radiography, in abdominal aortic aneurysm diagnosis, 851
Radionuclide scanning, for cholelithiasis detection, ultrasound versus, 825
in lower extremity deep vein thrombosis evaluation, 867–868
Rash, as travel vaccine reaction, 263–264
Reactive airway disease, ipratropium bromide for, 315–316
Recreation, for infants and children during travel, 76–77
Recreational boating injuries, 245
Rectal anticonvulsants, use with pediatric patient, 385
Rectal temperatures, 305
Red glass test, 661
Referrals, telemedicine services, 247
Reflexes, in neurologic examination, 501
Refractive error, 663
Refusal of care, by patient. See *Patient refusal.*
Regional nerve blocks, for wound care, 418
Reglan, for migraine, 408–409
Renal arteries, in ultrasound examination of aorta, 853, 855
Renal colic, diagnosis, 402–403
formal ultrasound evaluation, 878–881
pain management in, 405–406
ruptured abdominal aorta diagnosed as, 405
treatment, 403–405
Renal failure, formal ultrasound evaluation, 881
Renal infection, acute, formal ultrasound evaluation, 881–882
Renal stone, ultrasound image, 880
Renal ultrasound, **877–891**
emergency physicians performing, 884–891
formal, 878–884
acute renal infection, 881–882
acute renal vein thrombosis, 882
renal colic, 878–881
renal failure, 881
renal masses, 882
renal trauma, 882–884
Repatriation, injured marine crew members, 246–247
Reperfusion arrhythmias, 348
ACE inhibitors for, 348
Rescuers, at high altitude, 213
Resistance, to antimalarial drug, malaria parasite, 113–114
genetic, to malarial infection, 120–121
Resolution, ultrasound images, 937
axial. See *Axial resolution.*
beam width on, 948
lateral. See *Lateral resolution.*
Resperidone overdose, 443
Respiration, defining structures during ultrasound scan, 971
Respiratory care, pearls, autonomy for respiratory care practitioner, 315
ipratropium bromide, for reactive airways disease, 315–316
pitfalls, and updates, **315–325**
steroid metered-dose inhaler, for discharged chronic bronchospastics, 316
pitfalls, beta-agonists, for asthma patient, 319
cold agglutinin test, for *Mycoplasma pneumoniae,* 316–317
external chest compression, in severe asthma, 317
foreign body, in new-onset wheezers, 319–320
hypercarbia, correction in asthma and COPD, 318
initial ventilator settings, 317–318
prednisone, for discharged bronchospastic patient, 319
updates, antibiotics, choice for community-acquired pneumonia, 320–321
Bi-PAP versus CPAP, for acute pulmonary edema, 321–322
evaluation of suspected pulmonary embolism, 323–324
MDI-spacer therapy, for bronchospasm, 322–323
noninvasive ventilatory support, 321
permissive hypercapnia, 322
thrombolysis indications, in pulmonary embolism, 324–325
Respiratory care practitioner (RCP), autonomy for, 315
Respiratory distress, evaluation in pediatric patient, 382–383
Respiratory tract infections, in post-travel patients, 272–275
lung fluke infection, 273–274
melioidosis, 274
parasite-induced bronchospasm, 274–275
tuberculosis, 273
Rest, for pediatric migraine, 631
Resuscitation, central nervous system, **527–538**
calcium channel blockers and calpain inhibitors, 528–529
excitatory amino acid antagonists, 529–533
free radicals and lipid peroxidation, inhibitors of, 533–534
hypothermia, 534–535
inflammation inhibitors, 535–536
neurotrophic factors, 536–538
primary and secondary injury mechanisms, 527–528
cough CPR, 349

Resuscitation *(Continued)*
of hypothermic patient, 470
in prehospital setting, cessation of efforts, 297–298
fluid resuscitation, 296–297
in severe traumatic brain injury, algorithm for, 594
blood pressure, 590–592
spinal, in traumatic spinal cord injury, 708
Retinopathy, at high altitude, 205
Reverberation artifacts, ultrasound images, 963–964
foreign body detection, 897, 898
Reverse squeeze, 458
Rewarming techniques, for hypothermic patients, 470–472
active core, 470–472
active external, 470
passive, 470
Rhabdomyolysis, as poisoning complication, 436
Right ventricular dysfunction, echocardiographic diagnosis, 757
Riluzole, and glutamate release, 532–533
Risperdal (resperidone) overdose, 443
Rock foreign body, ultrasound image with shadow effect, 899
Romberg test, 502
Rotational acceleration injury, internal carotid artery and cervical spine relationship, 679
in mild head trauma, 564
Rotator cuff injury, ultrasound diagnosis, 376
Rotavirus, causing traveler's diarrhea, 182
Roundworms, in post-travel patients, 275, 276
Rubella immunization, pregnancy and, 98
routine, in childhood, 79
Rule-out ectopic pregnancy protocol, 800–803
Rupture, abdominal aorta aneurysm, misdiagnosis, 405
cardiac, echocardiographic diagnosis, 761
uterine, in pregnant trauma patient, ultrasound evaluation, 816
Ruptured ectopic pregnancy, sonographic appearance, 805, 806–807
tachycardia and, 389–390

Safety issues, ultrasound scanning, 972–974
foreign body detection, 902, 903–905
Sagittal section. See *Longitudinal scan, ultrasound.*
Sailing, medical emergencies and. See *Maritime medicine.*
St. Jude valves, 348
Saline, hypertonic, for blood pressure resuscitation, in severe traumatic brain injury, 592
for wound irrigation, 420
Salivary gland, masses in, 329
Salmonella, causing traveler's diarrhea, 182
SBIs. See *Secondary brain insults (SBIs).*
Scapular manipulation methods, of anterior shoulder dislocation, 365–366
Schistosomiasis, post-travel patient, 8–9, 268–269
Schmidt-Rimpler test, 504
Scombroid, 161, 162, 164
SCUBA diving, injury associated with. See *Diving-related emergencies.*
medical contraindications, 24
risks associated with, 23–25
Sea-faring vessels, medical emergencies on. See *Maritime medicine.*
Seated scapular manipulation method, anterior shoulder dislocation, 365
Secondary brain insults (SBIs), in severe traumatic brain injury, 582–584
outcome prediction data, 583, 584
Second trimester, vaginal bleeding in, ultrasound evaluation, 813–815
placental abruption, 814–815
placenta previa, 813–814
Sedation, pediatric patient, 385–386
for severe traumatic brain injury patient, 598
Seizures, characteristic findings in, 639–640
evaluation in neurologic examination, 493
past and family history in, 640–641
physical examination in, 641–642
postspell state, 640
syncope versus, physical and laboratory findings, 638, 639
witnessed in Emergency Department, 642
Seldane, causing long QT arrhythmias, 349
Selective serotonin reuptake inhibitors (SSRIs), overdose, 442
pharmacokinetics, 669
serotonin syndrome and, 665–666, 668–669
Self-contained underwater breathing apparatus (SCUBA). See *SCUBA diving.*
(S)-emopamil, in CNS resuscitation, 528
Sensitivity, of diagnostic tests, 303–304
Sensory function, in neurologic examination, 500–501
Septic arthritis, in pediatric patient, 374–375
Sequential array scanner, ultrasound probe, 931–934
Serotonin, migraines and, 629

Serotonin 5-HT agonist, for migraine, 407
adverse effects, 407–408
Serotonin syndrome, 438, 440–441, **665–673**
clinical presentation, 670–671
diagnostic criteria for, 670
differential diagnosis, 671–672
drug interactions, 667–670
incidence, 440, 665–666
management, 672–673
pathophysiology, 666–667
signs and symptoms, 440
Serzone (nefazodone) overdose, 443
Severe traumatic brain injury, **581–600**
airway control, oxygenation, and ventilation, 588–590
blood pressure resuscitation, 590–592
management guidelines, 581–582
physiologic concepts, 585–588
secondary brain insults, 582–584
therapeutic agents and techniques, 592–600
analgesia, 598
barbiturates, 600
CT scanning, prioritization of, 596–598
Glasgow Coma Scale, 593, 595
hyperventilation, 599–600
hypothermia, 600
management algorithm, 594
mannitol, 599
neuromuscular blockade, 598
patient monitoring, 595–596
sedation, 598
steroids, 600
transport protocol, 596–598
Shadowing, ultrasound images, acoustic, 944, 952–955, 956–959
lateral cystic, 960–961
Shaking, causing metaphyseal fractures in children, 372
Shear forces, in traumatic brain injury, 564
Shellfish poisoning, **157–175,** 163, 170. See also *Food poisoning.*
amnestic, 172
diarrheal, 172
neurologic, 171–172
paralytic, 171
treatment and prevention, 172
Shigella, causing traveler's diarrhea, 182
Shipping, medical emergencies and. See *Maritime medicine.*
Ship-to-shore communication, 247–248
Shock, in hemodynamically compromised patient, 431
Sickle cell disease, high altitude travel and, 209
vaso-occlusive crisis in, 400–401
patient-controlled analgesia for management of, 401–402
treatment inadequacy, 400–401
Side lobe artifacts, ultrasound images, 962–963
SIDS. See *Sudden infant death syndrome (SIDS).*
Sinus barotrauma, as diving-related emergency, 227–229
Sinus disorders, air travel and, 257
Sinuses, malignant lesions, 329
Sinus squeeze, 458
Sirens, use by emergency medical services vehicle, 291–293
Skin, decompression sickness affecting, 456
Skin lesions, in post-travel patients, 277–279
creeping eruption, 277–278
cutaneous leishmaniasis, 279
cutaneous myiasis, 278
Sleep disturbances, at high altitude, 194–195
treatment, 198
Sleeping pills, use by pregnant traveler, 103
Slit lamp examination, 660
SMA. See *Superior mesenteric artery (SMA).*
SMS. See *Subacute mountain sickness (SMS).*
Snakebites. See *Venomous snakebites (U.S.).*
Snellen chart, 660
SNX-111, in CNS resuscitation, 528
Society of Academic Emergency Physicians (SAEM), ultrasound training and credentialling guidelines, 914–915
SOD. See *Superoxide dismutase (SOD).*
Sodium channel blockade, in poisoned patient, 429–430
Sonodense structure, on ultrasound image, 940
diaphragm, 942
Sonolucent structure, on ultrasound image, 941
Sore throat, "normal," considerations for, 335–336
"strep," treatment of, 334–335
Sound waves, fluid- versus air-containing structures and, 922–923
frequency and amplitude, 924, 925. See also *Frequency.*
lateral cystic shadowing, 924, 925
and ultrasound, 922–924, 925. See also *Ultrasound images.*
Spacer, ultrasound scanning technique, 972
Specificity, of diagnostic tests, 303–304
Spells, cerebrovascular disease, 646
differential diagnosis and management strategies, **637–647**
diagnostic testing, 642–646
history, 639–641
physical examination, 641–642
drop attacks, 646
migraines, 646
syncope versus seizure, 638, 639

Spells *(Continued)*
transient global amnesia, 646
Spinal cord, anatomy, 699–701
Spinal cord dysfunction, as DCS manifestation, 230–231
Spinal cord syndromes, **699–709**
anatomical considerations, 699–701
anterior cord, 704
Brown-Sequard, 704
central cord, 704
complete cord, 703–704
conus medullaris, 705
management, 705–709
neurologic examination, 701–703
partial, 373–374
syringomyelia, 705
Spinal metastasis, 708–709
Spinal resuscitation, in traumatic spinal cord injury, 708
Spine, extremities and, **365–377**
Spontaneous dissections, carotid artery. See *Carotid artery dissection (CAD), spontaneous.*
vertebral artery. See *Vertebral artery dissections (VAD), spontaneous.*
Sporogony, 119
Sports activities, in pregnancy, 105–107
Sports injuries, and concussions, 575
management guidelines, 575
Spread of comitance, 663
SSRIs. See *Selective serotonin reuptake inhibitors (SSRIs).*
Stabilization, of patient before transfer, 287, 309–310
Stable arrhythmias, wide QRS rhythm, noninvasive diagnosis, 341–342
Staphylococcus aureus food poisoning, 6, 158–159
Static scanner, ultrasound probe, 936
Status epilepticus, pediatric, rectal anticonvulsants for, 385
Stereognosis, in sensory examination, 500
Steroids, for chronic bronchospastics, 316
for severe traumatic brain injury patient, 600
in treatment of pediatric meningitis, 385
for upper airway disease, 328
Stomach, ultrasound diagnosis of pyloric stenosis, 840–842
Strabismus, 663
Strep throat, treatment, 334–335
Streptococcal pharyngitis, treatment, 334–335
Streptococcus pneumonia isolates, pediatric meningitis and, 385
Streptokinase (STK), and iatrogenic compartment syndromes, 373
in treatment of acute myocardial infarction, 357–359
Stridor, in pediatric patients, and sore throat, case scenario, 335, 336
Stroke. See also *Stroke syndromes.*
incidence, 551
neuroimaging, 514–516
presenting signs and symptoms, 556
risks following therapy for acute myocardial infarction, 355
therapeutic window in, 551
in young people, 348
Stroke syndromes, **551–560**
anatomical considerations, 551–553
clinical presentation, 554–557
ischemic, 553
management, 557–560
pathophysiology, 553–554
Strongyloides, causing traveler's diarrhea, 183
Subacute mountain sickness (SMS), 207
Subarachnoid hemorrhage, cerebral computed axial tomography diagnosing, 304
misdiagnosis, 409–410
neuroimaging, 511, 512
Subcostal window, in 2-D echocardiography, 745–746, 747
in acute pulmonary embolism, 757–758
in cardiac injury diagnosis, 759–761
in cardiac tamponade diagnosis, 751–753, 755
Subdural hemorrhage, neuroimaging, 511–512
Subluxation, radial head, in children, 371–372
Submandibular gland tumors, 329
Succinylcholine, use in prehospital setting, 299
Suction, contraindicated for venomous snakebites, 474–475
Sudden infant death syndrome (SIDS), botulism and, 615
Sulfonamide, for malaria prevention in pregnant traveler, 100
Sumatriptan, for migraine, 407
for pediatric migraine, 631–632
serotonin syndrome and, 669–670
Sun, exposure to, at altitude, 206
as health care issue for travelers, 9–10
Sunscreens, 10
Superficial femoral vein (SFV), in ultrasound examination of lower extremity, 866
Superior mesenteric artery (SMA), in ultrasound examination of aorta, 853, 855
Superoxide dismutase (SOD), in CNS resuscitation, 533–534
Supine scapular manipulation method, anterior shoulder dislocation, 365–366
Supracondylar fractures, pediatric, failure to recognize, 366, 368

Supranuclear control, ocular motor system, 654
Suprapubic aspiration, ultrasound guidance for, 908
Supraventricular tachycardia, diagnosis, 345–346
Surgical procedures, ultrasound aiding, 742–743
Survivorship data, emergency medical services design and, 285–286
Suture line, aftercare and wound follow-up, 424
Suture techniques, for wound closure, 421–424
Swimming, as health care issues for travelers, 8–9
Sympathomimetic overdose, in serotonin syndrome differential diagnosis, 672
Syncope, during air travel, 255
  characteristic findings in, 639–640
  at high altitude, 205
  past and family history in, 640–641
  physical examination in, 641–642
  postspell state, 640
  versus seizure, physical and laboratory findings, 638, 639
  witnessed in Emergency Department, 642
Synovitis, toxic, in pediatric patient, 374–375
Syringe aspiration technique, confirming correct endotracheal intubation placement, 287
Syringomyelia, 705
Syrup of ipecac, 296

Tachycardia(s), pacemaker-mediated, 349–350
  ruptured ectopic pregnancy and, 389–390
Tachyphylaxis, 504
TAC (tetracaine, adrenaline, and cocaine) mixture, pediatric use, 386
Talk and deteriorate patients, minor head injury and, 568
Tandem gait, 502
Tapeworms, in post-travel patients, 275, 276
TAS. See *Transabdominal ultrasonography (TAS).*
TBI. See *Traumatic brain injury (TBI).*
TCAs. See *Tricyclic antidepressants (TCAs).*
TCDB. See *Traumatic Coma Data Bank (TCDB).*
Teardrop fractures, pediatric, failure to recognize, 366, 368
Telemedicine services, case management, 246
  communication logistics, 247–248
  education and training, 247
  evacuation and repatriation, 246–247
  medical advice, 245–246
  referrals, 247
Telephone, ship-to-shore communication, 247–248
Telex system, ship-to-shore communication, 248
Temazepam, for insomnia, with or without AMS, 201
Temperature(s), measurement in pediatric patient, 382
  rectal, 305
  tympanic, 304–305
Tensilon challenge test, 662
  in myasthenia gravis diagnosis, 614
Tension-type headache, ketorolac versus meperidine for, 408
Testicular torsion, diagnostic ultrasound for, 742
Tetanus, as cause of weakness, 610
Tetanus immunization, adult, 55
  routine, in childhood, 48, 78
  special groups recommendations, 66
Tetracaine, adrenaline, and cocaine (TAC) mixture, pediatric use, 386
Tetracycline, pregnant traveler and, diarrhea treatment, 108
  malaria prevention, 100
Tetrodotoxin (TTX), 168–170
TGC. See *Time gain compensator (TGC).*
Theophylline, atrioventricular block response to, 349
Therapeutic window, in stroke, 551, 557–558
Thermometers, tympanic, 304–305
Third trimester, vaginal bleeding in, ultrasound evaluation, 813–815
  placental abruption, 814–815
  placenta previa, 813–814
Threatened abortion, ultrasound evaluation, 811
Three arcs, carpal dislocations and, 365, 366
Thrombolytic therapy, for acute pulmonary embolism, 324–325
  compartment syndromes induced by, 373
  guidelines, 359
  versus percutaneous transluminal coronary angioplasty, for acute myocardial infarction, 354–357
  specific agents, 357–359. See also *Streptokinase (STK); Tissue plasminogen activator (tPA).*
  for stroke patient, 559
Thrombosis, during air travel, 258
  of cortical veins or venous sinus, 517–518

Thrombosis *(Continued)*
at high altitude, 204
ultrasound evaluation, acute renal vein, 882
central vein. See *Central vein thrombosis.*
deep vein. See *Deep vein thrombosis (DVT); Lower extremity deep vein thrombosis (LEDVT).*
Tickborne encephalitis immunization, 61
Tick paralysis, 610, 617
Tilt table test, in diagnosis of spells, 645–646
Time gain compensator (TGC), 943–944, 945
Tirilizad mesylate, in CNS resuscitation, 534
Tissue, changes at high altitude, 194
Tissue plasminogen activator (tPA), and iatrogenic compartment syndromes, 373
in treatment of acute myocardial infarction, 357–359
Tissue pressure, compartment syndromes and, 373
TNF. See *Tumor necrosis factor (TNF).*
TOA. See *Tubo-ovarian abscesses (TOA).*
Tonic water, causing long QT arrhythmias, 349
Topical anesthesia, for wound care, 418
Torsion, ovarian, sonographic findings, 820–821
Tourniquet examination, contraindicated for venomous snakebites, 474–475
of extremities, 371
Toxic hypoglycemic syndrome, 174
Toxicology, pearls, hazardous material incidents and psychologic injury, 433–434
hemodynamically compromised patient, approach to, 430–433
pitfalls, and updates in, **427–445**
poisoned patient evaluation, laboratory role in, 427–430
pitfalls, adverse drug reactions, recognizing, 437–438, 439
complications of poisoning, 435–436
serotonin syndrome, 438, 440–441
updates, *N*-acetylcysteine, emerging role of, 441–442
antidotes, 443–445
new psychotropic medicines, 442–443
Toxicology laboratory, role in evaluation of poisoned patient, 427–430
Toxic synovitis, in pediatric patient, 374–375
Toxin-induced weakness, 616
Clostridium botulinum, 615–616
Toxoplasma gondii infection, in AIDS patient, neuroimaging studies, 520–521, 522–524
Toxoplasmosis, in AIDS patient, neuroimaging studies, 520–521, 522–524
tPA. See *Tissue plasminogen activator (tPA).*
Traction headaches, 626–627
Training, in emergency ultrasound, 782, 784–785, 914–915
Tranquilizers, use by pregnant traveler, 103
Transabdominal ultrasonography (TAS), female pelvis, 789
versus endovaginal sonography, 790–792
normal anatomy, 792–793
Transducer, for ultrasound, annular array, 934
changing angle of, 971
contact scanner, 936
curved linear scanner, 934, 936
differences between, 936–937
electronic sector scanner, 934, 935
frequencies used, 937
positions, in blunt abdominal trauma evaluation, 766
principles, 930–931
rotating during scan, 970
in aorta examination, 854
types, linear array scanner, 931–934
mechanical sector scanner, 931, 932
Transesophageal echocardiography, update on, 359–361
Transfer of patients, 289
COBRA rules, 287–289
guidelines, 289
screening examination requirements, 309–310
Transferring hospital, responsibilities, 288
screening examination requirements, 309–310
Transient global amnesia, 646
Transient ischemic attack (TIA), 554
Transient unilateral monocular diplopia, 659, 660
Transthoracic echocardiography, update on, 359–361
Transverse scan, ultrasound, 738
kidney, 886, 887
pylorus, 841
Trauma. See also *Trauma patients.*
during air travel, 259
cardiac, echocardiographic diagnosis, 759–761
to head, classification, 565
incidence, 563
mild. See *Mild head trauma.*
severe. See *Resuscitation, central nervous system; Severe traumatic brain injury.*
nonaccidental fractures, 370, 372
to pregnant patient, pelvic ultrasound evaluation in, 816

while traveling, 96–98
pressure-related. See *Barotrauma.*
renal, formal ultrasound evaluation, 882–884
Trauma patients, hypothermia in, 472
transesophageal echocardiography in, 360–361
Traumatic brain injury (TBI), incidence, 563
mild. See *Mild head trauma.*
severe. See *Resuscitation, central nervous system; Severe traumatic brain injury.*
Traumatic Coma Data Bank (TCDB), outcome data in severe brain injury patients, 582–584
Traumatic spinal cord injury (TSCI), incidence, 706
initial assessment in, 706–708
pharmacologic therapy for, 708
spinal resuscitation in, 708
Travel, commercial air, medical emergencies in, **251–259.** See also *Commercial air travel.*
illness related to, epidemiology, 2
immunizations and, **43–69.** See also *Immunizations.*
with infants and children, **71–90.** See also *Infants and children, travel with.*
pregnancy and, **93–109.** See also *Pregnant traveler.*
wilderness, medical limitations to, **17–38**
Travelers, emergency department visits by, 261–280
diarrhea, 268–272
malaria, 264–267
respiratory tract infections, 272–275
skin lesions, 277–279
travel vaccine reactions, 261–264
worms and parasites, 275–277
general medical advice for, **1–14**
health care, 5–13
high altitude trips, 210–211, 221–222
insurance, 4–5
post-trip evaluation, 13–14
pre-trip planning, 3–4
immunization needs, 44, 45. See also *Immunizations.*
Traveler's diarrhea, **179–188**
causes, 179, 182–183
high-risk patients for, 25
high-risk versus low-risk destinations, 179
incidence, 179
in infants and children, 88–89
management, 181
in emergency department, 268
patient education, 184
treatment, 184–188
in pregnant traveler, 108–109
risk factors, 179–180
workup and diagnosis, 180–181
Travel insurance, 4–5
for pregnant traveler, 95–96
Travel kits. See *Medical kits.*
Travel medicine, establishment, 1
immunizations and, 43–44. See also *Immunizations.*
pre-trip evaluation, 2
problems encountered in emergency department, **261–280**
diarrhea, 268–272
malaria, 264–267
respiratory tract infections, 272–275
skin lesions, 277–279
vaccine reactions, 261–264
worms and parasites, 275–277
Travel vaccine reactions, 261–264
anaphylaxis or serious hypersensitivity reactions, 262
gastrointestinal symptoms, 262–263
local pain and swelling, 263
rash, fever, and associated symptoms, 263–264
Trazodone overdose, 443
Trekking, at high altitude, 210–211
information sources, 221–222
mountain sickness prevention, 199
by pregnant traveler, 107
Triage, for minors, 311
Trichinella, causing traveler's diarrhea, 183
Trichuriasis, causing traveler's diarrhea, 183
Tricyclic antidepressants (TCAs), overdose, 442
serotonin syndrome and, 665–666, 668, 669
Trigeminal nerve (cranial nerve V), in neurologic examination, 497
Trigger points, muscle spasm and, 618–619
Trip. See also *Travel entries.*
medical evaluation following, 13–14
medical planning prior to, 3–4
Trochlear nerve (cranial nerve IV), diplopia and, 653, 657
in neurologic examination, 496–497
Trochlear palsy, 657
L-Tryptophan, serotonin syndrome and, 665–666
TSCI. See *Traumatic spinal cord injury (TSCI).*
TTX. See *Tetrodotoxin (TTX).*
Tubal ring, ectopic pregnancy and, 804–805, 806
Tuberculosis, in post-travel patients, ED management, 273
pre-trip immunization, 64–65
for infants and children, 84
Tubo-ovarian abscesses (TOA), sonographic findings, 817, 818
Tumor necrosis factor (TNF), in CNS resuscitation, 535–536

Tumors, of the brain, neuroimaging, 523–524
Tungiasis, 8
Twiddler syndrome, 342
Two-point discrimination test, in sensory examination, 500
Tympanic membrane, barotrauma to, 226–227
  and temperature measurement in pediatric patient, 382
Tympanic temperatures, 304–305
Tympanostomy, for persistent effusions associated with otitis media, 328
Typhoid fever, in post-travel patient, ED management, 269
  pre-trip immunization, 55, 57–59
    for elderly, 66
    for immunocompromised person, 66
    for infants and children, 66, 84–85
    in pregnancy, 66, 99

UDS. See *Urine drug screen (UDS).*
Ultrasound, abdominal, **825–843.** See also *Abdominal ultrasound.*
  biologic effects and safety considerations, 972–974
  in blunt abdominal trauma, **763–785**
    in adults, 774–781
    in children, 781–782
    equipment, 765–766
    evaluation algorithm, 783
    history, 763–765
    limitations, 774
    technique, 766–774
    training, 782–785
  cardiac, **745–761.** See also *Echocardiography.*
  detection of foreign bodies and procedure guidance, **895–910**
    echo patterns, 897–901
    localization and retrieval, 901–902
    material and detection techniques, 895–897
    potential applications in emergency department, 902, 904–910
    safety issues, 902, 903–905
  effects and artifacts, 944, 946–965, 966–969
  emergency, credentialling issues in, **913–919**
    qualification pathway, 915–916
    quality assurance, 916–917
    strategies, 917–919
    training and guidelines, 914–915
  in the emergency department, applications of, **735–743**
    examination characteristics, 736–737
    future applications, 740–743
    limitations, 737–739
    scope, 739–740
  fundamentals, **921–974**
  images. See *Ultrasound images.*
  instrumentation, basic, 930–937
  kidney. See *Renal ultrasound.*
  lower extremity, 866. See also *Lower extremity deep vein thrombosis (LEDVT).*
  pelvic, **789–821.** See also *Pelvic ultrasound.*
  in penetrating extremity injury evaluation, 377
  pictures provided by, 924–930
  in placental separation diagnosis, 394–395
  renal, **877–891.** See also *Renal ultrasound.*
  in rotator cuff injury diagnosis, 376
  scanning techniques, 965, 969–972
  sound waves and, 922–924, 925
  upper extremity examination, 870–871. See also *Central vein thrombosis.*
  vascular, **849–871.** See also *Vascular ultrasound.*
Ultrasound images, adult polycystic kidney disease, 888
  aorta, 852–856
  appendix, 838
  cirrhosis of the liver, 943
  diaphragm, 942
  effects and artifacts, beam-width artifact, 961–962
    central effects, 944, 952–960
    common artifacts, 968
    lateral cystic shadowing, 960–961
    mirror effects, 965, 966–967, 969
    reverberation artifacts, 963–964
    side lobe artifacts, 962–963
  emphysematous cholecystitis, 958
  foreign bodies, 898–901
  gain, 941–944
  gallstones. See *Gallstones.*
  gray scale versus white, 939–942
  hydronephrosis, 880, 888–890
  intussusception, 843
  kidney, 880, 941–942
  liver, 835
  M mode echocardiography, 940
  qualities, 924, 926–930
    A- and B-modes, 926–927
    density of objects, 929
    depth and size of objects, 928, 930
  renal stone, 880
  resolution, 937
    axial and lateral, 944, 946–951
  translation conventions, 937–939
  ureterovesical junction, 890
  vesicoureteral reflux, 889
Universal precautions kit, commercial airline, 253
Unscheduled landings, in-flight medical emergencies, 254–255

Upper airway disease, infections in pediatric patient, 383
steroids for, 328
Upper extremity, ultrasound examination, 870–871
Ureterovesical junction, ultrasound image, 890
Urine drug screen (UDS), 430
Uterus, empty, ectopic pregnancy and, 809
ruptured, in pregnant trauma patient, 816
ultrasonography examination, endovaginal, 794–796
pregnant trauma patient, 816
transabdominal, 795

Vaccines. See also *Immunizations.*
adverse reactions to, 2, 261–264. See also *Travel vaccine reactions.*
Haemophilus influenzae (HIB), 383
during pregnancy, 27
for travelers, 26
VAD. See *Vertebral artery dissections (VAD).*
Vaginal bleeding, ectopic pregnancy and, 389
ultrasound evaluation, first trimester, 800–811. See also *Ectopic pregnancy.*
second- and third-trimester, 813–815
Vagus nerve (cranial nerve X), in neurologic examination, 499
Valsalva maneuver, ultrasound scanning technique, 972
Varicella immunization, adult, 55
routine, in childhood, 50, 80
special groups recommendations, 66
Vascular duplex scanning, in nontraumatic CNS dysfunction, 510
Vascular ultrasound, **849–871**
abdominal aortic aneurysm, 849–858
Doppler principles, 859–860
in emergency department, 859
peripheral, 860–868
deep vein thrombosis, central vein, 868–870, 871
isolated calf-vein, 868
lower extremity, 860–868
upper extremity and central veins, 870–871
Vasoconstrictors, for wound care, 418–419
Vasodilators, for HAPE, 203
Vaso-occlusive crisis (VOC), 400–401
patient-controlled analgesia for, 401–402
treatment inadequacy in, 400–401
Vehicle, emergency medical services, patient transfer by, 289
use of lights and siren by, 291–292
Veins, ultrasound scanning, localization procedure, 907
visualization technique, 907–908
Venlafaxine overdose, 443
Venomous snakebites (U.S.), 473–476
incidence, 473
pathophysiology, 474
pit vipers, 473–474
treatment, contraindicated, 474–475
indicated, 475–476
Venous access, in pediatric patient, 382
Ventilation, in sever traumatic brain injury, 588–590
Ventilation/perfusion (V/Q), in pediatric patient, 383
in suspected pulmonary embolism evaluation, 323, 324
Ventilatory changes, at high altitude, 193
Ventilatory support, noninvasive, 321
preventing volutrauma, 317–318
Ventricular tachycardia, diagnosis, 345–347
Venturi-type oxygen mask, limited use, 296
Verapamil, dysrhythmias and, diagnosis, 345, 346
Vergence, 654, 663
Versed (midazolam), for pediatric patient, 385–386
Vertebral artery, carotid artery and, dissections, **677–695.** See also *Carotid artery dissection (CAD); Vertebral artery dissections (VAD).*
diagnostic evaluation, 691–692
diagnostic pitfalls and dilemmas, 693–695
intracranial, 688
pathophysiology, 678–680
spontaneous, 680–688
traumatic, 688–691
treatment, 692–693
Vertebral artery dissections (VAD), pain distribution in, 685
spontaneous, 684–688
case report, 684
clinical course or prognosis, 687–688
clinical features, 686
clinical presentation, 684–685
intracranial, 688
predisposing factors and mechanisms of injury, 685, 687
traumatic, 688
mild blunt injury, 689
serious blunt or penetrating injuries, 690–691
Vertigo, vestibular nerve and, 498
Very hot intravenous fluids, hypothermic patient, 471
Vesicoureteral reflux, ultrasound image, 889
Vessels, ultrasound scanning, localization procedure, 907
visualization technique, 907–908

Vestibular decompression sickness, 231
Vestibular nerve, in neurologic examination, 498
VIBRATED SPASMS mnemonic, 705, 706
Vibration testing, in sensory examination, 500–501
Vibrio cholera, causing traveler's diarrhea, 182
Video conferencing, ship-to-shore communication, 248
Visual disturbances, evaluation in neurologic examination, 493
VOC. See *Vaso-occlusive crisis (VOC).*
Volutrauma, ventilatory management and, 317–318
V/Q. See *Ventilation/perfusion (V/Q).*

Water, precautions concerning, infants and children, 75–76
  international travel and, 6
  pregnant traveler, 108
Water purification, for diarrhea prevention, 184
  pregnant traveler, 105
Weakness, case presentations, 609, 612–613, 621–622
  causes, 610–611
  disorders presenting with, bites and ticks, 617
    botulism, 615–616
    controversial and poorly defined syndromes, 618–620
    drug- or toxin-induced weakness, 616
    electrolyte-induced weakness, 616–617
    fish poisoning, 617
    Guillain-Barré syndrome, 613
    Lambert-Eaton myasthenic syndrome, 615
    myasthenia gravis, 614–615
    psychiatric syndromes, 618–620
  evaluation, 605–609, 610–611
    algorithm for, 606–607
    drug use, 607, 608
    in neurologic examination, 493, 499–500
    patient history, 606–607
    signs and symptoms, 608–609
    testing protocols, 609
  facial nerve, 498
  generalized, and peripheral neuromuscular disease, approach to, **605–622**
  testing for, 499–500
    protocols, 609, 610–611
Weber's syndrome, 656, 664
Wheezing, in pediatric patients, foreign body and, 319–320
  and sore throat, case scenario, 335, 336
White structure, on ultrasound image, 940
  diaphragm, 942
Wilderness travel, medical limitations to, **17–38**
    coronary artery disease, 30–34
    diabetes, 34
    ear, nose, and throat disorders, 36
    elderly, 30
    environmental risks, 18–26. See also *Environmental risks.*
    gastrointestinal disorders, 35
    HIV infection and immunocompromised persons, 35
    infants and children, 28–30
    mental stability, 37
    musculoskeletal disorders, 35
    neurologic conditions, 36
    obesity, 36
    participant classification, 37
    physical disabilities, 35–36
    in pregnancy, 26–28
    pulmonary disease, 34
  organizational resources, 41
Wood foreign body, ultrasound image with shadow effect, 899
Worms, in post-travel patients, 275–277
  roundworms, 276
  tapeworms, 276–277
Wound care, pearls, pitfalls, and updates, **417–424**
    suture line aftercare and wound follow-up, 424
    wound closure, 421–424
  pitfalls, wound contamination, 419–420
    wound inspection, 420–421
  updates, anesthesia, 417–419

X-ray images, ultrasound images versus, 923–924

Yellow fever immunization, 55, 85–86
  country requirements, 60–61
  for elderly, 66
  geographic considerations, 62
  for immunocompromised person, 66
  for infants and children, 66
  in pregnancy, 66, 98–99
Yersinia, causing traveler's diarrhea, 182
Yolk sac, sonographic evaluation, 798